

Aidan Ryther

CSCI 761 - Topics In Advanced Algorithms

Assignment 3

Part 1:

Graph 2 Automorphism Groups Mappings:

- 0 1 2 3 4 5 6 7 -> 0 1 2 3 4 5 6 7
- 0 1 2 3 4 5 6 7 -> 0 1 3 2 4 6 5 7
- 0 1 2 3 4 5 6 7 -> 1 0 2 3 7 5 6 4
- 0 1 2 3 4 5 6 7 -> 1 0 3 2 7 6 5 4
- 0 1 2 3 4 5 6 7 -> 2 3 0 1 5 4 7 6
- 0 1 2 3 4 5 6 7 -> 2 3 1 0 5 7 4 6
- 0 1 2 3 4 5 6 7 -> 3 2 0 1 6 4 7 5
- 0 1 2 3 4 5 6 7 -> 3 2 1 0 6 7 4 5

Graph 2 Automorphism Groups Cycle Notation:

- ()
- (2 3)(5 6)
- (0 1)(4 7)
- (0 1)(2 3)(4 7)(5 6)
- (0 2)(1 3)(4 5)(6 7)
- (0 2 1 3)(4 5 7 6)
- (0 3 1 2)(4 6 7 5)
- (0 3)(1 2)(4 6)(5 7)

Graph 3 Automorphism Groups Mappings:

- 0 1 2 3 4 5 6 7 -> 0 1 2 3 4 5 6 7
- 0 1 2 3 4 5 6 7 -> 0 2 1 3 5 4 7 6
- 0 1 2 3 4 5 6 7 -> 1 0 4 6 2 7 3 5
- 0 1 2 3 4 5 6 7 -> 1 4 0 6 7 2 5 3
- 0 1 2 3 4 5 6 7 -> 2 0 5 7 1 6 3 4
- 0 1 2 3 4 5 6 7 -> 2 5 0 7 6 1 4 3
- 0 1 2 3 4 5 6 7 -> 3 6 7 0 5 4 1 2

- 0 1 2 3 4 5 6 7 -> 3 7 6 0 4 5 2 1
- 0 1 2 3 4 5 6 7 -> 4 1 7 5 0 3 6 2
- 0 1 2 3 4 5 6 7 -> 4 7 1 5 3 0 2 6
- 0 1 2 3 4 5 6 7 -> 5 2 6 4 0 3 7 1
- 0 1 2 3 4 5 6 7 -> 5 6 2 4 3 0 1 7
- 0 1 2 3 4 5 6 7 -> 6 3 5 1 7 2 0 4
- 0 1 2 3 4 5 6 7 -> 6 5 3 1 2 7 4 0
- 0 1 2 3 4 5 6 7 -> 7 3 4 2 6 1 0 5
- 0 1 2 3 4 5 6 7 -> 7 4 3 2 1 6 5 0

Graph 3 Automorphism Groups Cycle Notation:

- ()
- (1 2)(4 5)(6 7)
- (0 1)(2 4)(3 6)(5 7)
- (0 1 4 7 3 6 5 2)
- (0 2 5 6 3 7 4 1)
- (0 2)(1 5)(3 7)(4 6)
- (0 3)(1 6)(2 7)(4 5)
- (0 3)(1 7)(2 6)
- (0 4)(2 7)(3 5)
- (0 4 3 5)(1 7 6 2)
- (0 5 3 4)(1 2 6 7)
- (0 5)(1 6)(3 4)
- (0 6)(1 3)(2 5)(4 7)
- (0 6 4 2 3 1 5 7)
- (0 7 5 1 3 2 4 6)
- (0 7)(1 4)(2 3)(5 6)

Part 2:

The entries where I used the clause "if both $R(s,t-1)$ and $R(s-1,t)$ are even" are underlined.

	3	4	5	6	7	8	9	10
3	6	<u>9</u>	14	<u>19</u>	26	<u>33</u>	42	<u>51</u>
4		18	<u>31</u>	50	<u>75</u>	108	<u>149</u>	200
5			62	<u>111</u>	186	<u>293</u>	442	<u>641</u>
6				222	<u>407</u>	700	<u>1141</u>	1782
7					814	<u>1513</u>	2654	<u>4435</u>
8						3026	<u>5679</u>	10114
9							11358	<u>21471</u>
10								42942

The upper bounds on $R(s,s)$ implied by $R(s,s) \leq \text{choose}(2s-2, s-1)$ for $3 \leq s \leq 10$ are:

- $S = 3 \rightarrow \text{choose}(4,2) = 6$
- $S = 4 \rightarrow \text{choose}(6,3) = 20$
- $S = 5 \rightarrow \text{choose}(8,4) = 70$
- $S = 6 \rightarrow \text{choose}(10,5) = 252$
- $S = 7 \rightarrow \text{choose}(12,6) = 924$
- $S = 8 \rightarrow \text{choose}(14,7) = 3432$
- $S = 9 \rightarrow \text{choose}(16,8) = 12870$
- $S = 10 \rightarrow \text{choose}(18,9) = 48620$

The the upper bounds on $R(s,s)$ for $3 \leq s \leq 10$ are:

- $S = 3 \rightarrow R(3,3) \leq 55$
- $S = 4 \rightarrow R(4,4) \leq 209$
- $S = 5 \rightarrow R(5,5) \leq 793$
- $S = 6 \rightarrow R(6,6) \leq 3011$
- $S = 7 \rightarrow R(7,7) \leq 11442$
- $S = 8 \rightarrow R(8,8) \leq 43478$
- $S = 9 \rightarrow R(9,9) \leq 165217$

- $S = 10 \rightarrow R(10,10) \leq 627822$

Part 3:

The lower bounds on $R(s,s)$ for $3 \leq s \leq 10$ are:

- $S = 3 \rightarrow R(3,3) \geq 3$
- $S = 4 \rightarrow R(4,4) \geq 5$
- $S = 5 \rightarrow R(5,5) \geq 6$
- $S = 6 \rightarrow R(6,6) \geq 9$
- $S = 7 \rightarrow R(7,7) \geq 12$
- $S = 8 \rightarrow R(8,8) \geq 17$
- $S = 9 \rightarrow R(9,9) \geq 23$
- $S = 10 \rightarrow R(10,10) \geq 33$

The best which can be obtained numerically from $2^{\binom{n}{s}} / 2^{\binom{s}{2}} < 1$ are:

- $S = 3 \rightarrow n = 3 \rightarrow R(3,3) \geq 4$
- $S = 4 \rightarrow n = 6 \rightarrow R(4,4) \geq 7$
- $S = 5 \rightarrow n = 11 \rightarrow R(5,5) \geq 12$
- $S = 6 \rightarrow n = 17 \rightarrow R(6,6) \geq 18$
- $S = 7 \rightarrow n = 27 \rightarrow R(7,7) \geq 28$
- $S = 8 \rightarrow n = 42 \rightarrow R(8,8) \geq 43$
- $S = 9 \rightarrow n = 65 \rightarrow R(9,9) \geq 66$
- $S = 10 \rightarrow n = 100 \rightarrow R(10,10) \geq 101$