

Small Ramsey Numbers

Stanisław P. Radziszowski
Department of Computer Science
Rochester Institute of Technology
Rochester, NY 14623, spr@cs.rit.edu
<http://www.cs.rit.edu/~spr>

Submitted: June 11, 1994; Accepted: July 3, 1994
Revision #12: August 4, 2009

ABSTRACT: We present data which, to the best of our knowledge, includes all known nontrivial values and bounds for specific graph, hypergraph and multicolor Ramsey numbers, where the avoided graphs are complete or complete without one edge. Many results pertaining to other more studied cases are also presented. We give references to all cited bounds and values, as well as to previous similar compilations. We do not attempt complete coverage of asymptotic behavior of Ramsey numbers, but concentrate on their specific values.

Mathematical Reviews Subject Number 05C55.

Revisions

1993, February	preliminary version, RIT-TR-93-009 [Ra2]
1994, July 3	accepted to the <i>EJJC</i> , posted on the web
1994, November 7	<i>EJJC</i> revision #1
1995, August 28	<i>EJJC</i> revision #2
1996, March 25	<i>EJJC</i> revision #3
1997, July 11	<i>EJJC</i> revision #4
1998, July 9	<i>EJJC</i> revision #5
1999, July 5	<i>EJJC</i> revision #6
2000, July 25	<i>EJJC</i> revision #7
2001, July 12	<i>EJJC</i> revision #8
2002, July 15	<i>EJJC</i> revision #9
2004, July 4	<i>EJJC</i> revision #10
2006, August 1	<i>EJJC</i> revision #11
2009, August 4	<i>EJJC</i> revision #12

Table of Contents

1. Scope and Notation	3
2. Classical Two Color Ramsey Numbers	4
2.1 Bounds on $R(k, l)$ for $k \leq 10, l \leq 15$	4
2.2 Bounds on $R(k, l)$ for $l \geq 15$	6
2.3 Other results on $R(k, l)$	7
3. Two Colors - Most Studied Cases	9
3.1 Dropping one edge from complete graph	9
3.2 Complete bipartite graphs	11
3.3 Cycles, cycles versus complete graphs	14
3.4 Cycles versus wheels	16
3.5 Cycles versus books	17
4. General Graph Numbers in Two Colors	18
4.1 Paths	18
4.2 Wheels	18
4.3 Books	19
4.4 Trees and forests	19
4.5 Stars, stars versus other graphs	20
4.6 Fans, fans versus other graphs	21
4.7 Paths versus other graphs	21
4.8 Triangle versus other graphs	21
4.9 Cycles versus other graphs	22
4.10 Wheels versus other graphs	23
4.11 Books versus other graphs	23
4.12 Trees and forests versus other graphs	24
4.13 Cases for $n(G), n(H) \leq 5$	24
4.14 Mixed cases	25
4.15 Multiple copies of graphs, disconnected graphs	26
4.16 General results for sparse graphs	26
4.17 Other general two color results	27
5. Multicolor Ramsey Numbers	29
5.1 Bounds for classical numbers	29
5.2 General results for complete graphs	31
5.3 Cycles	32
5.4 Paths, paths versus other graphs	35
5.5 Special cases	36
5.6 Other general results	36
6. Hypergraph Numbers	38
7. Cumulative Data and Surveys	40
7.1 Cumulative data for two colors	40
7.2 Cumulative data for three colors	41
7.3 Surveys	41
8. Concluding Remarks	42
References	44-72

1. Scope and Notation

There is vast literature on Ramsey type problems starting in 1930 with the original paper of Ramsey [Ram]. Graham, Rothschild and Spencer in their book [GRS] present an exciting development of Ramsey Theory. The subject has grown amazingly, in particular with regard to asymptotic bounds for various types of Ramsey numbers (see the survey papers [GrRö, Neš, ChGra2]), but the progress on evaluating the basic numbers themselves has been unsatisfactory for a long time. In the last three decades, however, considerable progress has been obtained in this area, mostly by employing computer algorithms. The few known exact values and several bounds for different numbers are scattered among many technical papers. This compilation is a fast source of references for the best results known for specific numbers. It is not supposed to serve as a source of definitions or theorems, but these can be easily accessed via the references gathered here.

Ramsey Theory studies conditions when a combinatorial object contains necessarily some smaller given objects. The role of Ramsey numbers is to quantify some of the general existential theorems in Ramsey Theory.

Let G_1, G_2, \dots, G_m be graphs or s -uniform hypergraphs (s is the number of vertices in each edge). $R(G_1, G_2, \dots, G_m; s)$ denotes the m -color **Ramsey number** for s -uniform graphs/hypergraphs, avoiding G_i in color i for $1 \leq i \leq m$. It is defined as the least integer n such that, in any coloring with m colors of the s -subsets of a set of n elements, for some i the s -subsets of color i contain a sub-(hyper)graph isomorphic to G_i (not necessarily induced). The value of $R(G_1, G_2, \dots, G_m; s)$ is fixed under permutations of the first m arguments. If $s=2$ (standard graphs) then s can be omitted. If G_i is a complete graph K_k , then we can write k instead of G_i , and if $G_i = G$ for all i we can use the abbreviation $R_m(G; s)$ or $R_m(G)$. For $s=2$, $K_k - e$ denotes a K_k without one edge, and for $s=3$, $K_k - t$ denotes a K_k without one triangle (hyperedge).

The graph nG is formed by n disjoint copies of G , and the **join** $G+H$ of vertex disjoint graphs G and H is obtained by adding all the edges between vertices of G and H . P_i is a **path** on i vertices, C_i is a **cycle** of length i , and W_i is a **wheel** with $i-1$ spokes, i.e. a graph formed by some vertex x , connected to all vertices of some cycle C_{i-1} (thus $W_i = K_1 + C_{i-1}$). $K_{n,m}$ is a complete n by m bipartite graph, in particular $K_{1,n}$ is a **star** graph. The **book** graph $B_i = K_2 + \bar{K}_i = K_1 + K_{1,i}$ has $i+2$ vertices, and can be seen as i triangular pages attached to a single edge. The **fan** graph F_n is defined by $F_n = K_1 + nK_2$. For a graph G , $n(G)$ and $e(G)$ denote the number of vertices and edges, respectively, and $\delta(G)$ and $\Delta(G)$ minimum and maximum degree in G . Finally, let $\chi(G)$ be the chromatic number of G . In general we follow the notation used by West [West].

Section 2 contains the data for the classical two color Ramsey numbers $R(k, l)$ for complete graphs, and section 3 for the most studied two color cases. Section 4 lists other often studied two color cases for general graphs. The multicolor and hypergraph cases are gathered in sections 5 and 6, respectively. Finally, section 7 gives pointers to cumulative data and to the other surveys.

2. Classical Two Color Ramsey Numbers

2.1. Upper and lower bounds on $R(k, l)$

l	3	4	5	6	7	8	9	10	11	12	13	14	15	
k														
3	6	9	14	18	23	28	36	40 43	46 51	52 59	59 69	66 78	73 88	
4		18	25	35 41	49 61	56 84	73 115	92 149	97 191	128 238	133 291	141 349	153 417	
5			43 49	58 87	80 143	101 216	125 316	143 442	159 633	185 848	209 1139	235 1461	265 1878	
6				102 165	113 298	130 495	169 780	179 1171	253 1804	262 2566	317 3705		401 6911	
7					205 540	216 1031	237 1713	289 2826	405 4553	416 6954	511 10581		22116	
8						282 1870	317 3583		6090	10630	16944	27490	817 41525	861 63620
9							565 6588	580 12677		22325	39025	64871	89203	
10								798 23556			81200			1265

Table I. Known nontrivial values and bounds for two color Ramsey numbers $R(k, l) = R(k, l; 2)$.

l	4	5	6	7	8	9	10	11	12	13	14	15
k												
3	GG	GG	Kéry	Ka2 GY	GR MZ	Ka2 GR	Ex5 RK2	Ka2 RK2	Ex12 Les	Piw1 RK2	Ex8 RK2	WW Les
4	GG	Ka1 MR4	Ex9 MR5	Ex3 Mac	Ex15 Mac	Ex17 Mac	HaKr Mac	2.3.e Spe3	SLL Spe3	2.3.e Spe3	XXR Spe3	XXR Spe3
5		Ex4 MR5	Ex9 HZ1	CET Spe3	HaKr Spe3	Ex17 Mac	Ex17 Mac	Ex17 HW+	Ex17 HW+	Ex17 HW+	Ex17 HW+	Ex17 HW+
6			Ka1 Mac	Ex17 Mac	XSR2 Mac	XXER Mac	Ex17 Mac	XXR HW+	2.3.e HW+	XXER HW+		2.3.h HW+
7				She1 Mac	2.3.e Mac	XSR2 HZ1	2.3.h Mac	XXER HW+	2.3.e HW+	XXR HW+		HW+
8					BR Mac	XXER Ea1	HZ1	HW+	HW+	XXER HW+		2.3.h HW+
9						She1 ShZ1	2.3.e Ea1	HW+	HW+	HW+		
10							She1 Shi2			Yang		2.3.h

References for Table I. HW+ abbreviates HWSYZH.

We split the data into the table of values and a table with corresponding references. In Table I, known exact values appear as centered entries, lower bounds as top entries, and upper bounds as bottom entries. For some of the exact values two references are given when the lower and upper bound credits are different.

The task of proving $R(3, 3) \leq 6$ was the second problem in Part I of the William Lowell Putnam Mathematical Competition held in March 1953 [Bush]. Greenwood and Gleason [GG] in 1955 established the initial values $R(3,4)=9$, $R(3,5)=14$ and $R(4,4)=18$. Kéry [Kéry] in 1964 found $R(3,6)=18$, but only recently an elementary and self-contained proof of this result appeared in English [Car].

All the critical graphs for the numbers $R(k, l)$ (graphs on $R(k, l) - 1$ vertices without K_k and without K_l in the complement) are known for $k = 3$ and $l = 3, 4, 5$ [Kéry], 6 [Ka2], 7 [RK3, MZ], and there are 1, 3, 1, 7 and 191 of them, respectively. All $(3, k)$ -graphs, for $k \leq 6$, were enumerated in [RK3], and all $(4, 4)$ -graphs in [MR2]. There exists a unique critical graph for $R(4, 4)$ [Ka2]. There are 430215 such graphs known for $R(3, 8)$ [McK], 1 for $R(3, 9)$ [Ka2] and 350904 for $R(4, 5)$ [MR4], but there might be more of them. In [MR5] evidence is given for the conjecture that $R(5, 5) = 43$ and that there exist 656 critical graphs on 42 vertices. The graphs constructed by Exoo in [Ex9, Ex12, Ex13, Ex14, Ex15, Ex16, Ex17], and some others, are available electronically from <http://ginger.indstate.edu/ge/RAMSEY>.

The construction by Mathon [Mat] and Shearer [She1] (see also sections 2.3.i, 5.2.h and 5.2.i), using data obtained by Shearer [She3] for primes up to 7000, gives the following lower bounds for higher diagonal numbers: $R(11, 11) \geq 1597$, $R(13, 13) \geq 2557$, $R(14, 14) \geq 2989$, $R(15, 15) \geq 5485$, and $R(16, 16) \geq 5605$. Similarly, $R(17, 17) \geq 8917$, $R(18, 18) \geq 11005$ and $R(19, 19) \geq 17885$ were obtained in [LSL], though the first two of these bounds follow also from the data in [She3]. The same approach does not improve on an easy bound $R(12, 12) \geq 1637$ [XXR], which can be obtained by applying twice 2.3.e. Only some of the higher bounds implied by 2.3.* are shown, and more similar bounds could be easily derived. In general, we show bounds beyond the contiguous small values if they improve on results previously reported in this survey or published elsewhere. Some easy upper bounds implied by 2.3.a are marked as [Ea1].

Cyclic (or *circular*) graphs are often used for Ramsey graph constructions. Several cyclic graphs establishing lower bounds were given in the Ph.D. dissertation by J.G. Kalbfleisch in 1966, and many others were published in the next few decades (see [RK1]). Only recently Harborth and Krause [HaKr] presented all best lower bounds up to 102 from cyclic graphs avoiding complete graphs. In particular, no lower bound in Table I can be improved with a cyclic graph on less than 102 vertices. See also item 2.3.k and section 4.16 [HaKr].

The claim that $R(5, 5) = 50$ posted on the web [Stone] is in error, and despite being shown to be incorrect more than once, this value is still being cited by some authors. The bound $R(3, 13) \geq 60$ [XieZ] cited in the 1995 version of this survey was shown to be incorrect in [Piw1]. Another incorrect construction for $R(3, 10) \geq 41$ was described in [DuHu].

There are really only two general upper bound inequalities useful for small parameters, namely 2.3.a and 2.3.b. Stronger upper bounds for specific parameters were difficult to obtain, and they often involved massive computations, like those for the cases of (3,8) [MZ], (4,5) [MR4], (4,6) and (5,5) [MR5]. The bound $R(6, 6) \leq 166$, only 1 more than the best known [Mac], is an easy consequence of a theorem in [Walk] (2.3.b) and $R(4, 6) \leq 41$. T. Spencer [Spe3], Mackey [Mac], and Huang and Zhang [HZ1], using the bounds for minimum and maximum number of edges in (4,5) Ramsey graphs listed in [MR3, MR5], were able to establish new upper bounds for several higher Ramsey numbers, improving on all of the previous longstanding results by Giraud [Gi3, Gi5, Gi6].

We have recomputed the upper bounds in Table I marked [HZ1] using the method from the paper [HZ1], because the bounds there relied on an overly optimistic personal communication from T. Spencer. Further refinements of this method are studied in [HZ2, ShZ1, Shi2]. The paper [Shi2] subsumes the main results of the manuscripts [ShZ1, Shi2]. The upper bound marked in Table I [Yang] was obtained by Yang using the method of [HWSYZH] (abbreviated in the table as HW+).

2.2. Bounds on $R(k, l)$, higher parameters

l	15	16	17	18	19	20	21	22	23
k									
3	73 WW	79 WW	92 WWY1	99 Ex17	106 WWY1	111 Ex17	122 WWY1	131 WSLX2	137 WSLX2
4	153 XXR	163 Ex17	182 LSS1	187 2.3.e	213 2.3.g	234 Ex17	242 SLZL	282 SL	
5	265 Ex17	289 2.3.h	388 XSR2	395 2.3.e	407 XSR2	421 2.3.h	441 2.3.h	485 2.3.h	521 2.3.h
6	401 2.3.h	434 SLLL	548 SLLL	614 SLLL	710 SLLL	878 SLLL		1070 SLLL	
7			711 2.3.g	725 2.3.h	908 SLLL		1214 SLLL		
8	861 2.3.h		937 ShaXP	1045 2.3.g	1236 2.3.g		1617 2.3.h		

l	24	25	26	27	28	29	30	31
k								
3	143 WSLX1	154 WSLX2	159 WSLX1	167 WSLX1	173 WSLX2	184 WSLX2	190 WSLX2	199 WSLX2

l	32	33	34	35	36	37	38	39
k								
3	214 WSLX2	218 WuCXS	226 WuCXS	231 WuCXS	239 WuCXS	244 WuCXS	256 WuCXS	

Table II. Known nontrivial lower bounds for higher two color Ramsey numbers $R(k, l)$, with references.

The upper bounds of 88, 99, 110, 121, 133, 145, 158 on $R(3, k)$ for $15 \leq k \leq 21$, respectively, were obtained in [Les]. The lower bounds marked [XXR], [XXER], 2.3.e and 2.3.h need not be cyclic. Several of the Cayley colorings from [Ex17] are also non-cyclic. All other lower bounds listed in Table II were obtained by construction of cyclic graphs. The graphs establishing lower bounds marked 2.3.g can be constructed by using appropriately chosen graphs G and H with a common m -vertex induced subgraph, similarly as it was done in several cases in [XXR].

Yu [Yu2] constructed a special class of triangle-free cyclic graphs establishing several lower bounds for $R(3, k)$, for $k \geq 61$. Only one of these bounds, $R(3, 61) \geq 479$, cannot be easily improved by the inequality $R(3, 4k+1) \geq 6R(3, k+1) - 5$ from [CCD] (2.3.c) and data from Tables I and II. Finally, for higher parameters we mention two more cases which improve on bounds listed in earlier revisions: $R(9, 17) \geq 1411$ is given in [XXR] and $R(10, 15) \geq 1265$ can be obtained by using 2.3.h.

In general, one can expect that the lower bounds in Table II are weaker than those in Table I, in the sense that with some work many of them should not be hard to improve, in contrast to the bounds in Table I, especially smaller ones.

2.3. Other results on $R(k, l)$

- (a) $R(k, l) \leq R(k-1, l) + R(k, l-1)$, with strict inequality when both terms on the right hand side are even [GG]. There are obvious generalizations of this inequality for avoiding graphs other than complete.
- (b) $R(k, k) \leq 4R(k, k-2) + 2$ [Walk].
- (c) Explicit construction for $R(3, 4k+1) \geq 6R(3, k+1) - 5$, for all $k \geq 1$ [CCD].
- (d) Constructive results on triangle-free graphs in relation to the case of $R(3, k)$ [BBH1, BBH2, Fra1, Fra2, FrLo, Gri, KM1, Loc, RK3, RK4, Stat, Yu1].
- (e) Bounds for the difference between consecutive Ramsey numbers, in particular the bound $R(k, l) \geq R(k, l-1) + 2k - 3$ for $k, l \geq 3$ [BEFS].
- (f) By taking a disjoint union of two critical graphs one can easily see that $R(k, p) \geq s$ and $R(k, q) \geq t$ imply $R(k, p+q-1) \geq s+t-1$. Xu and Xie [XX1] improved this construction to yield better general lower bounds, in particular $R(k, p+q-1) \geq s+t+k-3$.
- (g) For $2 \leq p \leq q$ and $3 \leq k$, if (k, p) -graph G and (k, q) -graph H have a common induced subgraph on m vertices without K_{k-1} , then $R(k, p+q-1) > n(G) + n(H) + m$. In particular, this implies the bounds $R(k, p+q-1) \geq R(k, p) + R(k, q) + k - 3$ and $R(k, p+q-1) \geq R(k, p) + R(k, q) + p - 2$ [XX1, XXR].
- (h) $R(2k-1, l) \geq 4R(k, l-1) - 3$ for $l \geq 5$ and $k \geq 2$, and in particular for $k=3$ we obtain $R(5, l) \geq 4R(3, l-1) - 3$ [XXER].
- (i) If the quadratic residues Paley graph Q_p of prime order $p = 4t+1$ contains no K_k , then $R(k, k) \geq p+1$ and $R(k+1, k+1) \geq 2p+3$ [She1, Mat]. Data for larger p was obtained in [LSL]. See also 3.1.c, and items 5.2.h and 5.2.i for similar multicolor results.

- (j) Study of Ramsey numbers for large disjoint unions of graphs [Bu1, Bu9], in particular $R(nK_k, nK_l) = n(k+l-1) + R(K_{k-1}, K_{l-1}) - 2$, for n large enough [Bu8].
- (k) $R(k, l) \geq L(k, l) + 1$, where $L(k, l)$ is the maximal order of any cyclic (k, l) -graph. A compilation of many best cyclic bounds was presented in [HaKr].
- (l) The graphs critical for $R(k, l)$ are $k-1$ vertex connected and $2k-4$ edge connected, for $k, l \geq 3$ [BePi].
- (m) Two color lower bounds can be obtained by using items 5.2.k, 5.2.l and 5.2.m with $r = 2$. Some generalizations of these were obtained in [ZLLS].

In the last six items of this section we only briefly mention some pointers to the literature dealing with asymptotics of Ramsey numbers. This survey was designed mostly for small, finite, and combinatorial results, but still we wish to give the reader some useful and representative references to more traditional papers looking first of all at the infinite.

- (n) In a 1995 breakthrough Kim proved that $R(3, k) = \Theta(k^2/\log k)$ [Kim].
- (o) Explicit triangle-free graphs with independence k on $\Omega(k^{3/2})$ vertices [Alon2, CPR].
- (p) Other general and asymptotic results on triangle-free graphs in relation to the case of $R(3, k)$ [AKS, Alon2, CCD, CPR, Gri, FrLo, Loc, She2].
- (q) In 1947, Erdős gave an amazingly simple probabilistic proof that $R(k, k) \geq c \cdot k 2^{k/2}$ [Erd1]. Spencer [Spe1] improved the constant in the last result. More probabilistic asymptotic lower bounds for other Ramsey numbers were obtained in [Spe1, Spe2, AlPu].
- (r) Other asymptotic bounds for $R(k, k)$ can be found, for example, in [Chu3, McS] (lower bound) and [Tho, Con1] (upper bound), and for many other bounds in the general case of $R(k, l)$ consult [Spe2, GRS, GrRö, Chu4, ChGra2, LiRZ1, AlPu, Kriv].
- (s) Explicit construction of a graph with clique and independence k on $2^c \log^2 k / \log \log k$ vertices by Frankl and Wilson [FraWi]. Further constructions by Chung [Chu3] and Grolmusz [Grol1, Grol2]. Explicit constructions like these are usually weaker than known probabilistic results.

3. Two Colors - Most Studied Cases

3.1. Dropping one edge from complete graph

This section contains known values and nontrivial bounds for the two color case when the avoided graphs are complete or have the form $K_k - e$, but not both are complete.

G	H	$K_3 - e$	$K_4 - e$	$K_5 - e$	$K_6 - e$	$K_7 - e$	$K_8 - e$	$K_9 - e$	$K_{10} - e$	$K_{11} - e$
$K_3 - e$		3	5	7	9	11	13	15	17	19
K_3		5	7	11	17	21	25	31	37 38	42 47
$K_4 - e$		5	10	13	17	28	29 38	34	41	
K_4		7	11	19	27 36	37 52				
$K_5 - e$		7	13	22	31 39	40 66				
K_5		9	16	30 34	43 67	112				
$K_6 - e$		9	17	31 39	45 70	59 135				
K_6		11	21	37 55	116	205				
$K_7 - e$		11	28	40 66	59 135	251				
K_7		13	28 34	51 88	202					

Table III. Two types of Ramsey numbers $R(G, H)$, includes all known nontrivial values.

- (a) The exact values in Table III involving $K_3 - e$ are obvious, since one can easily see that $R(K_3 - e, K_k) = R(K_3 - e, K_{k+1} - e) = 2k - 1$, for all $k \geq 2$.
- (b) The bound $R(K_4 - e, K_8) \leq 45$ is given in [LinMac], and $R(K_3, K_{12} - e) \geq 46$ in [MPR]. Wang, Wang and Yan [WWY2] constructed cyclic graphs showing $R(K_3, K_{13} - e) \geq 54$, $R(K_3, K_{14} - e) \geq 59$ and $R(K_3, K_{15} - e) \geq 69$. It is known that $R(K_4, K_{12} - e) \geq 128$ [Shao] using one color of the (4,4,4;127)-coloring defined in [HiIr].
- (c) If the quadratic residues Paley graph Q_p of prime order $p = 4t + 1$ contains no $K_k - e$, then $R(K_{k+1} - e, K_{k+1} - e) \geq 2p + 1$. In particular, $R(K_{14} - e, K_{14} - e) \geq 2987$ [LiShen]. See also item 2.3.i.

G	H	K_4-e	K_5-e	K_6-e	K_7-e	K_8-e	K_9-e	$K_{10}-e$	$K_{11}-e$
K_3		CH2	Clan	FRS1	GH	Ra1	Ra1	MPR MPR	WWY2 MPR
K_4-e		CH1	FRS2	McR	McR	Ea1 HZ2	Ex14	Ex14	
K_4		CH2	EHM1	Ex11 Ea1	Ex14 HZ2				
K_5-e		FRS2	CEHMS	Ex14 Ea1	Ex14 HZ2				
K_5		BH	Ex8 Ex8	Ea1 HZ2	HZ2				
K_6-e		McR	Ex14 Ea1	Ex14 HZ2	Ex14 HZ2				
K_6		McN	Ex14 Ea1	HYZ	ShZ2				
K_7-e		McR	Ex14 HZ2	Ex14 HZ2	ShZ1				
K_7		Ea1 Ea1	Ex14 ShZ2	HYZ					

References for Table III.

- (d) More bounds (not shown in Table III) can be obtained by using Table I, an obvious generalization of the inequality $R(k, l) \leq R(k-1, l) + R(k, l-1)$, and by monotonicity of Ramsey numbers, in this case $R(K_{k-1}, G) \leq R(K_k - e, G) \leq R(K_k, G)$.
- (e) All $(K_3, K_l - e)$ -graphs for $l \leq 6$ have been enumerated [Ra1].
- (f) For the following the critical graphs are unique: $R(K_3, K_l - e)$ for $l=3$ [Tr], 6 and 7 [Ra1], $R(K_4 - e, K_4 - e)$ [FRS2], $R(K_5 - e, K_5 - e)$ [Ra3] and $R(K_4 - e, K_7 - e)$ [McR].
- (g) The number of $R(K_3, K_l - e)$ -critical graphs for $l = 4, 5$ and 8 is 4, 2 and 9, respectively [MPR], and there are at least 6 such graphs for $R(K_3, K_9 - e)$ [Ra1].
- (h) All the critical graphs for the cases $R(K_4 - e, K_4)$ [EHM1], $R(K_4 - e, K_5)$ and $R(K_5 - e, K_4)$ [DzFi1] are known, and there are 5, 13 and 6 of them, respectively.
- (i) $R(K_k - e, K_k - e) \leq 4R(K_{k-2}, K_k - e) - 2$ [LiShen].
For a similar inequality for complete graphs see 2.3.b.
- (j) The upper bounds from [ShZ1, ShZ2] are subsumed by a later article [Shi2].
- (k) The upper bounds in [HZ2] were obtained by a reasoning generalizing the bounds for classical numbers in [HZ1]. Several other results from section 2.3 apply, though checking in which situation they do may require looking inside the proofs whether they still hold for $K_n - e$.

3.2. Complete bipartite graphs

NOTE: This subsection gathers information on Ramsey numbers where specific bipartite graphs are avoided in a coloring of K_n (as everywhere in this survey), in contrast to often studied bipartite Ramsey numbers (not covered in this survey) where the initial coloring is of a bipartite graph $K_{n,m}$.

Numbers

The following Tables IVa and IVb gather information mostly from the surveys by Lortz and Mengersen [LoM3, LoM4]. All cases involving $K_{1,2} = P_3$ are solved by a formula for $R(P_3, G)$, holding for all isolate-free graphs G , derived in [CH2]. All star versus star numbers are given by 3.2.a.

m, n	p, q	1, 2	1, 3	1, 4	1, 5	1, 6	2, 2	2, 3	2, 4	2, 5	3, 3	3, 4
2, 2		4 CH2	6 CH2	7 Par3	8 Par3	9 FRS4	6 CH1					
2, 3		5 CH2	7 FRS4	9 Stev	10 FRS4	11 FRS4	8 HaMe4	10 Bu4				
2, 4		6 CH2	8 HaMe3	9 Stev	11 HaMe4	13 LoM4	9 HaMe4	12 ExRe	14 EHM2			
2, 5		7 CH2	9 HaMe3	11 Stev	13 Stev	14 LoM4	11 HaMe4	13 LoM3	16 LoM1	18 EHM2		
2, 6		8 CH2	10 HaMe3	11 Stev	14 Stev	15* Shao	12 HaMe4	14 LoM3	17 LoM3	20 LoM1		
3, 3		7 CH2	8 HaMe3	11 LoM4	12 LoM4	13 LoM4	11 Lortz	13 HaMe3	16 LoM4	18 LoM4	18 HaMe3	
3, 4		7 CH2	9 HaMe3	11 LoM4	13 LoM4	14 LoM4	11 Lortz	14 LoM4	17 Sh+	≤21 LoM4	≤25 LoM2	≤30 LoM2
3, 5		9 CH2	10 HaMe3	13 Sh+	15 Sh+		14 HaMe4	≥15* Shao	≥16* Shao	≥21* Shao	≤28 LoM2	≤33 LoM2

Table IVa. Ramsey numbers $R(K_{m,n}, K_{p,q})$.
(unpublished results are marked with a *, Sh+ abbreviates ShaXBP)

- (a) The next few easily computed values of $R(K_{1,n}, K_{2,2})$, extending data in the first row of Table IVa, are 13, 14, 21 and 22 for n equal to 9, 10, 16 and 17, respectively. See function $f(n)$ in 3.2.c of the next subsection below.
- (b) Formula for $R(K_{1,n}, K_{k_1, k_2}, \dots, k_t, m)$ for m large enough, in particular for $t=1, k_1=2$ with $n \leq 5, m \geq 3$ and $n=6, m \geq 11$, for example $R(K_{1,5}, K_{2,7}) = 15$ [Stev].

m	2	3	4	5	6	7	8	9	10	11
6	12 HaMe4	14 LoM3	17 LoM3	20 LoM1	21 EHM2					
7	14 HaMe4	17 LoM3	19 LoM3	21 LoM3	24 LoM1	26 EMH2				
8	15 HaMe4	18 LoM3	20 LoM3	22*-23 LoM3	24-25 LoM3	28 LoM1	30 EMH2			
9	16 HaMe4	19 LoM3	22 LoM3	25* Shao	27* Shao	29* Shao	32 LoM1	33 EHM2		
10	17 HaMe4	21 LoM3	24 LoM3	27 LoM3	27-29 LoM3	28-31 LoM3	32-33 LoM3	36 LoM1	38 EHM2	
11	18 HaMe4						≤35 LoM3	36-37 LoM3	40 LoM1	42 EHM2

Table IVb. Known Ramsey numbers $R(K_{2,n}, K_{2,m})$, for $6 \leq n \leq 11$, $2 \leq m \leq 11$.
(unpublished results are marked with a *, Sh+ abbreviates ShaXBP)

- (c) The values and bounds for higher cases of $R(K_{2,2}, K_{2,n})$ are 20, 22, 22/23, 22/24, 25, 26, 27/28, 28/29, 30 and 32 for $12 \leq n \leq 21$, respectively. More exact values can be found for prime powers $\lceil \sqrt{n} \rceil$ and $\lceil \sqrt{n} \rceil + 1$ [HaMe4].
- (d) The known values of $R(K_{2,2}, K_{3,n})$ are 15, 16, 17, 20 and 22 for $6 \leq n \leq 10$ [Lortz], and $R(K_{2,2}, K_{3,11}) = 24$ [Shao]. See Tables IVa and IVb for the smaller cases, and [HaMe4] for upper bounds and values for some prime powers $\lceil \sqrt{n} \rceil$.
- (e) $R(K_{2,n}, K_{2,n})$ is equal to 46, 50, 54, 57 and 62 for $12 \leq n \leq 16$, respectively. The first open diagonal case is $65 \leq R(K_{2,17}, K_{2,17}) \leq 66$ [EHM2]. The status of all higher cases for $n < 30$ is listed in [LoM1].
- (f) $R(K_{1,4}, K_{4,4}) = R(K_{1,5}, K_{4,4}) = 13$ [ShaXPB]
 $R(K_{1,4}, K_{1,2,3}) = R(K_{1,4}, K_{2,2,2}) = 11$ [GuSL]
 $R(K_{1,7}, K_{2,3}) = 13$ [Par4]
 $R(K_{1,15}, K_{2,2}) = 20$ [La2]
 $R(K_{2,2}, K_{4,4}) = 14$ [HaMe4]
 $R(K_{2,2}, K_{4,5}) = 15$ [Shao]
 $R(K_{2,2}, K_{4,6}) = 16$ [Shao]
 $R(K_{2,2}, K_{5,5}) = R(K_{2,3}, K_{3,5}) = 17$ [Shao]
 $R(K_{3,5}, K_{3,5}) \leq 38$ [LoM2]
 $R(K_{4,4}, K_{4,4}) \leq 62$ [LoM2]

General results

- (a) $R(K_{1,n}, K_{1,m}) = n + m - \varepsilon$, where $\varepsilon = 1$ if both n and m are even and $\varepsilon = 0$ otherwise [Har1]. It is also a special case of multicolor numbers for stars obtained in [BuRo1].
- (b) $R(K_{1,3}, K_{m,n}) = m + n + 2$ for $m, n \geq 1$ [HaMe3].
- (c) $R(K_{1,n}, K_{2,2}) = f(n) \leq n + \sqrt{n} + 1$, with $f(q^2) = q^2 + q + 1$ and $f(q^2 + 1) = q^2 + q + 2$ for every q which is a prime power [Par3]. Furthermore, $f(n) \geq n + \sqrt{n} - 6n^{11/40}$ [BEFRS5]. For more bounds and values of $f(n)$ see [Par5, Chen, ChenJ, MoCa].
- (d) $R(K_{1,n+1}, K_{2,2}) \leq R(K_{1,n}, K_{2,2}) + 2$ [Chen].
- (e) $R(K_{2,\lambda+1}, K_{1,\nu-k+1})$ is either $\nu + 1$ or $\nu + 2$ if there exists a (ν, k, λ) -difference set. This and other related results are presented in [Par4, Par5]. See also [GoCM, GuLi].
- (f) Formulas and bounds on $R(K_{2,2}, K_{2,n})$, and bounds on $R(K_{2,2}, K_{m,n})$. In particular, $R(K_{2,2}, K_{2,k}) = n + k\sqrt{n} + c$, for $k = 2, 3, 4$ and some prime powers $\lceil \sqrt{n} \rceil$ and $\lceil \sqrt{n} \rceil + 1$, for some $-1 \leq c \leq 3$ [HaMe4].
- (g) $R(K_{2,n}, K_{2,n}) \leq 4n - 2$ for all $n \geq 2$, and the equality holds iff a strongly regular $(4n - 3, 2n - 2, n - 2, n - 1)$ -graph exists [EHM2].
- (h) Conjecture that $4n - 3 \leq R(K_{2,n}, K_{2,n}) \leq 4n - 2$ for all $n \geq 2$. Many special cases are solved and several others are discussed in [LoM1].
- (i) $R(K_{2,n-1}, K_{2,n}) \leq 4n - 4$ for all $n \geq 3$, with the equality if there exists a symmetric Hadamard matrix of order $4n - 4$. There are only 4 cases in which the equality does not hold for $3 \leq n \leq 58$, namely 30, 40, 44 and 48 [LoM1].
- (j) $R(K_{2,n-s}, K_{2,n}) \leq 4n - 2s - 3$ for $s \geq 2$ and $n \geq s + 2$, with the equality in many cases involving Hadamard matrices or strongly regular graphs. Asymptotics of $R(K_{2,n}, K_{2,m})$ for $m \gg n$ [LoM3].
- (k) Some algebraic lower and upper bounds on $R(K_{s,n}, K_{t,m})$ for various combinations of n , m and $1 \leq t, s \leq 3$ [BaiLi, BaLX].
- (l) Upper bounds for $R(K_{2,2}, K_{m,n})$ for $m, n \geq 2$, with several cases identified for which the equality holds. Special focus on the cases for $m = 2$ [HaMe4].
- (m) Bounds for the numbers of the form $R(K_{k,n}, K_{k,m})$, specially for fixed k and close to the diagonal cases. Asymptotics of $R(K_{3,n}, K_{3,m})$ for $m \gg n$ [LoM2].
- (n) $R(nK_{1,3}, mK_{1,3}) = 4n + m - 1$ for $n \geq m \geq 1$, $n \geq 2$ [BES].
- (o) Asymptotics for $K_{2,m}$ versus K_n [CLRZ].
- (p) Upper bound asymptotics for $K_{k,m}$ versus K_n [LiZa1].
- (q) Special two-color case applies in the study of asymptotics for multicolor Ramsey numbers for complete bipartite graphs [ChGra1].

3.3. Cycles, cycles versus complete graphs

Cycles

$$R(C_3, C_3) = 6 \text{ [GG, Bush]}$$

$$R(C_4, C_4) = 6 \text{ [CH1]}$$

$$R(C_3, C_n) = 2n - 1 \text{ for } n \geq 4 \text{ [ChaS]}$$

Result obtained independently in [Ros1] and [FS1], a new simpler proof in [KáRos]:

$$R(C_n, C_m) = \begin{cases} 2n - 1 & \text{for } 3 \leq m \leq n, m \text{ odd, } (n, m) \neq (3, 3) \\ n - 1 + m/2 & \text{for } 4 \leq m \leq n, m \text{ and } n \text{ even, } (n, m) \neq (4, 4) \\ \max \{ n - 1 + m/2, 2m - 1 \} & \text{for } 4 \leq m < n, m \text{ even and } n \text{ odd} \end{cases}$$

$$R(nC_3, mC_3) = 3n + 2m \text{ for } n \geq m \geq 1, n \geq 2 \text{ [BES]}$$

$$R(nC_4, mC_4) = 2n + 4m - 1 \text{ for } m \geq n \geq 1, (n, m) \neq (1, 1) \text{ [LiWa1]}$$

Formulas for $R(nC_4, mC_5)$ [LiWa2]

Formulas and bounds for $R(nC_m, nC_m)$ [Den, Biel1]

Unions of cycles, formulas and bounds for various cases including diagonal, different lengths, different multiplicities [MiSa, Den], and their relation to 2-local Ramsey numbers [Biel1].

Cycles versus complete graphs

Since 1976, it was conjectured that $R(C_n, K_m) = (n - 1)(m - 1) + 1$ for all $n \geq m \geq 3$, except $n = m = 3$ [FS4, EFRS2]. The parts of this conjecture were proved as follows: for $n \geq m^2 - 2$ [BoEr], for $n > 3 = m$ [ChaS], for $n \geq 4 = m$ [YHZ1], for $n \geq 5 = m$ [BJYHRZ], for $n \geq 6 = m$ [Schi1], for $n \geq m \geq 7$ with $n \geq m(m - 2)$ [Schi1], for $n \geq 7 = m$ [ChenCZ1], and for $n \geq 4m + 2, m \geq 3$ [Nik]. Open conjectured cases are marked in Table V by "conj."

	C_3	C_4	C_5	C_6	C_7	C_8	... C_n for $n \geq m$
K_3	6 GG-Bush	7 ChaS	9 ChaS	11 ChaS	13 ChaS	15 ChaS	... $2n-1$ ChaS
K_4	9 GG	10 CH2	13 He2/JR4	16 JR2	19 YHZ1	22 YHZ1	... $3n-2$ YHZ1
K_5	14 GG	14 Clan	17 He2/JR4	21 JR2	25 YHZ2	29 BJYHRZ	... $4n-3$ BJYHRZ
K_6	18 Kéry	18 Ex2-RoJa1	21 JR5	26 Schi1	31 Schi1	36 Schi1	... $5n-4$ Schi1
K_7	23 Ka2-GY	22 RT-JR1	25 Schi2	31 CheCZN	37 CheCZN	43 JarBa/Ch+	... $6n-5$ ChenCZ1
K_8	28 GR-MZ	26 RT		36 ChenCX	43 ChenCZ1	50 JarAl/ZZ3	... $7n-6$ conj.
K_9	36 Ka2-GR	30-32 RT-XSR1					... $8n-7$ conj.
K_{10}	40-43 Ex5-RK2	34-39 RT-XSR1					... $9n-8$ conj.

Table V. Known Ramsey numbers $R(C_n, K_m)$.
(Ch+ abbreviates ChenCZ1, comments on joint credits in (b) of this section)

- (a) The first column in Table V gives data from the first row in Table I.
- (b) Joint credit [He2/JR4] in Table V refers to two cases in which Hendry [He2] announced the values without presenting the proofs, which later were given in [JR4]. The special cases of $R(C_6, K_5) = 21$ [JR2] and $R(C_7, K_5) = 25$ were solved independently in [YHZ2] and [BJYHRZ]. The double pointer [JarBa/ChenCZ1] refers to two independent papers, similarly as [JarAl/ZZ3], except that in the latter case [ZZ3] refers to an unpublished manuscript. For joint credits marked in Table V with "-", the first reference is for the lower bound and the second for the upper bound.
- (c) Lower bound asymptotics [Spe2, FS4, AIRö].
- (d) Upper bound asymptotics [BoEr, FS4, EFRS2, CLRZ, Sud1, LiZa2, AIRö].
- (e) For cycles versus graphs other than complete see sections 3.4, 3.5 and 4.9.

3.4. Cycles versus wheels

Note: In this survey the wheel graph $W_n = K_1 + C_{n-1}$ has n vertices, while some authors use the definition $W_n = K_1 + C_n$ with $n + 1$ vertices. For the cases $W_3 = C_3$ versus C_m see section 3.3.

	C_3	C_4	C_5	C_6	C_7	C_8	C_m	for
W_4	9 GG	10 CH2	13 He2	16 JR2	19 YHZ1	22 ...	$3m - 2$...	$m \geq 4$ YHZ1
W_5	11 Clan	9 Clan	9 He4	11 JR2	13 SuBB2	15 ...	$2m - 1$...	$m \geq 5$ SuBB2
W_6	11 BE3	10 JR3	13 ChvS	16 SuBB2	19 ...	22 ...	$3m - 2$...	$m \geq 4$ SuBB2
W_7	13 BE3	9 Tse1				15*	$2m - 1^*$...	$m \geq 8$ ChenCMN
W_8	15 BE3	11 Tse1			19* ChenCN	22* ...	$3m - 2^*$...	$m \geq 7$ ChenCN
W_9	17 BE3	12 Tse1					$2m - 1^*$ ChenCMN	$m \geq 11$ ChenCMN
W_n for	... $2n - 1$ $n \geq 6$ BE3		$2n - 1$ $n \geq 19$ Zhou2		$2n - 1$ $n \geq 29$ Zhou2			cycles large wheels

Table VI. Ramsey numbers $R(W_n, C_m)$, for $n \leq 9, m \leq 8$. (results from unpublished manuscript are marked with a *)

- (a) $R(C_3, W_n) = 2n - 1$ for $n \geq 6$ [BE3]. All critical graphs have been enumerated. The critical graphs are unique for $n = 3, 5$, and for no other n [RaJi].
- (b) $R(C_4, W_n) = 13, 14, 16, 17$ for $n = 10, 11, 12, 13$, respectively [Tse1]
 $R(C_4, W_n) \leq n + \lceil (n - 1)/3 \rceil$ for $n \geq 7$ [SuBUB]
- (c) $R(W_n, C_m) = 2n - 1$ for odd m with $n \geq 5m - 6$ [Zhou2]
- (d) $R(W_n, C_m) = 3m - 2$ for even $n \geq 4$ with $m \geq n - 1, m \neq 3$, was conjectured by Surahmat et al. [SuBT1, SuBT2, Sur]. Parts of this conjecture were proved in [SuBT1, ZhaCC], and the proof was completed in [ChenCN].
- (e) Conjecture that $R(W_n, C_m) = 2m - 1$ for odd $n \geq 3$ and all $m \geq 5$ with $m > n$ [Sur]. It was proved for $2m \geq 5n - 7$ [SuBT1], and further for $2m \geq 3n - 5$ [ChenCMN].
- (f) Cycles are Ramsey unsaturated for some wheels [AliSur]
- (g) Study of cycles versus generalized wheels $W_{k,n}$ [Sur, SuBTB]

3.5. Cycles versus books

	C_3	C_4	C_5	C_6	C_7	C_8	C_9	C_{10}	C_{11}	C_m	for
B_2	7 RS1	7 <i>Fal6</i>	9 <i>Cal</i>	11 <i>Fal8</i>	13 ...	15	17	19	21	$2m - 1$...	$m \geq 2$ <i>Fal8</i>
B_3	9 RS1	9 <i>Fal6</i>	10 <i>Fal8</i>	11 JR2	13 <i>13</i>	15 <i>Fal8</i>	17 ...	19	21	$2m - 1$...	$m \geq 8$ <i>Fal8</i>
B_4	11 RS1	11 <i>Fal6</i>	11 <i>Fal8</i>	12	13	15	17	19 <i>Fal8</i>	21 ...	$2m - 1$...	$m \geq 10$ <i>Fal8</i>
B_5	13 RS1	12 <i>Fal6</i>	13 <i>Fal8</i>	14	15	15	17	19	21	$2m - 1$	$m \geq 12$ <i>Fal8</i>
B_6	15 RS1	13 <i>Fal6</i>	15 <i>Fal8</i>	16	17	18	18		21	$2m - 1$	$m \geq 14$ <i>Fal8</i>
B_7	17 RS1	16 <i>Fal6</i>	17 <i>Fal8</i>	16	19	20	21			$2m - 1$	$m \geq 16$ <i>Fal8</i>
B_8	19 RS1	17 Tse1	19 <i>Fal8</i>	17	19	22	≥ 23			$2m - 1$	$m \geq 18$ <i>Fal8</i>
B_9	21 RS1	18 Tse1	21 <i>Fal8</i>	18			≥ 25	≥ 26		$2m - 1$	$m \geq 20$ <i>Fal8</i>
B_{10}	23 RS1	19 Tse1	23 <i>Fal8</i>	19				≥ 28		$2m - 1$	$m \geq 22$ <i>Fal8</i>
B_{11}	25 RS1	20 Tse1	25 <i>Fal8</i>							$2m - 1$	$m \geq 24$ <i>Fal8</i>
B_n for	... $2n + 3$ $n \geq 2$ RS1	$\approx n$ some (c)	... $2n + 3$ $n \geq 4$ <i>Fal8</i>		$2n + 3$ $n \geq 15$ <i>Fal8</i>		$2n + 3$ $n \geq 23$ <i>Fal8</i>		$2n + 3$ $n \geq 31$ <i>Fal8</i>		cycles large books

Table VII. Ramsey numbers $R(B_n, C_m)$ for $n, m \leq 11$.
(using *et al.* abbreviations *Fal* for FRS and *Cal* for CRSPS)

- (a) For the cases of $B_1 = K_3$ versus C_m see section 3.3.
Centered entries in italics (middle of Table VII) are from personal communication and manuscripts by Shao. The latter include $R(B_n, C_n) \geq 3n - 2$, $R(B_{n-1}, C_n) \geq 3n - 4$ for $n \geq 3$, and an improvement to the bound on m in (e) to $m \geq 2n - 1 \geq 7$ [Shao].
- (b) $R(C_4, B_{12}) = 21$ [Tse1], $R(C_4, B_{13}) = 22$, $R(C_4, B_{14}) = 24$ [Tse2].
 $R(C_4, B_8) = 17$ [Tse2] (it was reported incorrectly in [FRS6] to be 16).
- (c) $q^2 + q + 2 \leq R(C_4, B_{q^2 - q + 1}) \leq q^2 + q + 4$ for prime power q [FRS6]. B_n is a subgraph of B_{n+1} , hence likely $R(C_4, B_n) = n + O(\sqrt{n})$ (compare to $R(C_4, K_{2,n})$ in section 3.2).
- (d) $R(B_n, C_m) = 2n + 3$ for odd $m \geq 5$ with $n \geq 4m - 13$ [FRS8]
- (e) $R(B_n, C_m) = 2m - 1$ for $n \geq 1$, $m \geq 2n + 2$ [FRS8]
- (f) More theorems on $R(B_n, C_m)$ in [FRS6, FRS8, NiRo4, Zhou1]

4. General Graph Numbers in Two Colors

This section includes data with respect to general graph results. We tried to include all nontrivial values and identities regarding exact results (or references to them), but only those out of general bounds and other results which, in our opinion, have a direct connection to the evaluation of specific numbers. If some small value cannot be found below, it may be covered by the cumulative data gathered in section 7, or be a special case of a general result listed in this section. Note that $P_2 = K_2$, $B_1 = F_1 = C_3 = W_3 = K_3$, $B_2 = K_4 - e$, $P_3 = K_3 - e$, $W_4 = K_4$ and $C_4 = K_{2,2}$ imply other identities not mentioned explicitly.

4.1. Paths

$$R(P_n, P_m) = n + \lfloor m/2 \rfloor - 1 \quad \text{for all } n \geq m \geq 2 \quad [\text{GeGy}]$$

Stripes mP_2 [CL1, CL2, Lor]

Disjoint unions of paths (also called linear forests) [BuRo2, FS2]

4.2. Wheels

Note: In this survey the wheel graph $W_n = K_1 + C_{n-1}$ has n vertices, while some authors use the definition $W_n = K_1 + C_n$ with $n + 1$ vertices.

m	3	4	5	6	7
n					
3	6	9 GG	11 Clan	11 BE3	13 BE3
4		18 GG	17 He3	19 FM	
5			15 He2	17 FM	
6				17 FM	

Table VIII. Ramsey numbers $R(W_n, W_m)$, for $n \leq m \leq 7$.

- (a) $R(W_3, W_n) = 2n - 1$ for all $n \geq 6$ [BE3]
All critical colorings for $R(W_3, W_n)$ for all $n \geq 3$ [RaJi]
- (b) The value $R(W_5, W_5) = 15$ was given in the Hendry's table [He2] without a proof. Later the proof was published in [HaMe2].
- (c) All critical colorings (2, 1 and 2) for $R(W_n, W_6)$ for $n = 4, 5, 6$ [FM]
- (d) $R(W_6, W_6) = 17$, $R(4,4) = 18$ and $\chi(W_6) = 4$ give a counterexample $G = W_6$ to the Erdős conjecture (see [GRS]) that $R(G, G) \geq R(K_{\chi(G)}, K_{\chi(G)})$.

4.3. Books

m	1	2	3	4	5	6	7
n							
1	6	7 CH2	9 Clan	11 RS1	13 RS1	15 RS1	17 RS1
2		10 CHI	11 Clan	13 Rou	16 RS1	17-18 Rou	≤ 20 FRS7
3			14 RS1	15 Sh+	17 RS1		
4				18 RS1	≤ 20 RS1	22 RS1	
5					21 RS1		
6						26 RS1	

Table IX. Ramsey numbers $R(B_n, B_m)$, for $n, m \leq 7$.
(Sh+ abbreviates ShaXBP)

- (a) $254 \leq R(B_{37}, B_{88}) \leq 255$ [Par6]
- (b) Recent computations established that $R(B_2, B_6) \leq 17$ (and thus equal to 17) and $R(B_2, B_7) = 18$, with 3 and 65 critical graphs, respectively [BLR].
- (c) $R(B_1, B_n) = 2n + 3$ for all $n > 1$ [RS1]
- (d) $R(B_n, B_m) = 2n + 3$ for all $n \geq cm$ for some $c < 10^6$ [NiRo2, NiRo3]
- (e) $R(B_n, B_n) = (4 + o(1))n$ [RS1, NiRS]
- (f) In general, $R(B_n, B_n) = 4n + 2$ for $4n + 1$ a prime power. Several other specific values (like $R(B_{62}, B_{65}) = 256$) and general equalities and bounds for $R(B_n, B_m)$ can be found in [RS1, FRS7, Par6, NiRS, LiRZ2].

4.4. Trees and forests

In this subsection T_n and F_n denote n -vertex tree and forest, respectively.

- (a) $R(T_n, T_n) \leq 4n + 1$ [EG]
- (b) $R(T_n, T_n) \geq \lfloor (4n - 1)/3 \rfloor$ [BE2], see also section 4.17
- (c) Conjecture that $R(T_n, T_n) \leq 2n - 2$, note that this is almost the same as asking if $R(T_n, T_n) \leq R(K_{1, n-1}, K_{1, n-1})$ [BE2], see also [Bu7, FSS1, ChGra2]. Discussion of the conjecture that $R(T_n, T_m) \leq n + m - 2$ holds for all trees [FSS1].

- (d) If $\Delta(T_n) = n - 2$, $\Delta(T_m) = m - 2$ then the exact values of $R(T_n, T_m)$ are known, and they are between $n + m - 5$ and $n + m - 3$ depending on n and m . In particular, for $n = 2k + 1$ $R(T_{2k+1}, T_{2k+1}) = 2n - 5$ [GuoV].
- (e) View tree T as a bipartite graph with parts t_1 and t_2 , $t_2 \geq t_1$. Define $b(T) = \max\{2t_1 + t_2 - 1, 2t_2 - 1\}$. Then the bound $R(T, T) \geq b(T)$ holds always, $R(T, T) = b(T)$ holds for many classes of trees [EFRS3, GeGy], and asymptotically [HaLT], but cases for nonequality have been found [GHK].
- (f) $R(F_n, F_n) > n + \log_2 n - O(\log \log n)$ [BE2], forests are tight for this bound [CsKo].
- (g) Comment in [BaLS] about a recent result of the authors of [AKS], which implies that $R(T_n, T_n) \leq 2n - 2$ holds for sufficiently large n .
- (h) $R(T_m, K_{1,n}) \leq m + n - 1$, with equality for $(m - 1) \mid (n - 1)$ [Bu1].
- (i) $R(T_m, K_{1,n}) = m + n - 1$ for sufficiently large n for almost all trees T_m [Bu1]. Many cases were identified for which $R(T_m, K_{1,n}) = m + n - 2$ [Coc, ZZ1], see also [Bu1].
- (j) $R(T_m, K_{1,n}) \leq m + n$ if T_n is not a star and $(m - 1) \nmid (n - 1)$, some classes of trees and stars for which the equality holds [GuoV].
- (k) Forests, linear forests (unions of paths) [BuRo2, FS3, CsKo].
- (l) Paths versus trees [FSS1], see also other parts of this survey involving special graphs, in particular sections 4.5, 4.7 and 4.12.

4.5. Stars, stars versus other graphs

$R(K_{1,n}, K_{1,m}) = n + m - \varepsilon$, where $\varepsilon = 1$ for even n and m , and $\varepsilon = 0$ otherwise [Har1]. This is also a special case of multicolor numbers for stars obtained in [BuRo1].

$R(K_{1,n}, K_m) = n(m - 1) + 1$ by Chvátal's theorem [Chv].

Stars versus C_4 [Par3, Par4, Par5, BEFRS5, Chen, ChenJ, GoMC, MoCa]

Stars versus $K_{2,n}$ [Par4, GoMC]

Stars versus $K_{n,m}$ [Stev, Par3]

Stars versus complete bipartite graphs [Par4, Stev]

See also section 3.2

$R(K_{1,4}, B_4) = 11$ [RS2]

$R(K_{1,4}, K_{1,2,3}) = R(K_{1,4}, K_{2,2,2}) = 11$ [GuSL]

$nK_{1,m}$ versus W_5 [BaHA]

Stars versus W_5 and W_6 [SuBa1]

Stars versus W_9 [Zhang2, ZhaCZ1]

Stars versus wheels [HaBA1, ChenZZ2, Kor]

Stars versus paths [Par2, BEFRS2]

Stars versus cycles [La1, Clark, see Par6]

Stars versus books [CRSPS, RS2]

Stars versus trees [Bu1, Cheng, Coc, GuoV, ZZ1]

Stars versus stripes mP_2 [CL1, CL2, Lor]

Stars versus $K_n - tK_2$ [Hua1, Hua2]
 Stars versus $2K_2$ [MO]
 Union of two stars [Gros2]
 Unions of stars versus wheels [BaHA, HaBA2]

4.6. Fans, fans versus other graphs

$R(F_1, F_n) = R(K_3, F_n) = 4n + 1$ for $n \geq 2$, and bounds for $R(F_m, F_n)$ [LR2, GGS]
 $R(F_2, F_n) = 4n + 1$ for $n \geq 2$ and $R(F_m, F_n) \leq 4n + 2m$ for $n \geq m \geq 2$ [LinLi]
 $R(K_4, F_n) = 6n + 1$ for $n \geq 3$ [SuBB3]
 Fans versus paths, formulas for a number of cases including $R(P_6, F_n)$ [SaBr2].
 Missing case $R(P_6, F_4) = 12$ solved in [Shao].
 Fans versus K_n [LR2]
 Lower bounds on $R(F_2, K_n)$ from cyclic graphs for $n \leq 9$ [Shao]

4.7. Paths versus other graphs

P_3 versus all isolate-free graphs [CH2]
 Paths versus stars [Par2, BEFRS2]
 Paths versus trees [FS4, FSS1]
 Paths versus books [RS2]
 Paths versus cycles [FLPS, BEFRS2]
 Paths versus K_n [Par1]
 Paths versus $K_{n,m}$ [Häg]
 Paths versus W_5 and W_6 [SuBa1]
 Paths versus W_7 and W_8 [Bas]
 Paths versus wheels [BaSu, ChenZZ1, SaBr3, Zhang1]
 Paths versus fans [SaBr2]
 Paths versus $K_1 + P_m$ [SaBr1, SaBr4]
 Paths and cycles versus trees [FSS1]
 Paths and unions of paths versus Jahangir graphs [AliBas, AliBT, AliSur]
 Sparse graphs versus paths and cycles [BEFRS2]
 Graphs with long tails [Bu2, BG]
 Monotone paths and cycles [Lef]
 Unions of paths [BuRo2]

4.8. Triangle versus other graphs

$R(3, k) = \Theta(k^2/\log k)$ [Kim]
 Explicit construction for $R(3, 4k + 1) \geq 6R(3, k + 1) - 5$, for all $k \geq 1$ [CCD]
 Explicit triangle-free graphs with independence k on $\Omega(k^{3/2})$ vertices [Alon2, CPR]
 $R(K_3, K_7 - 2P_2) = R(K_3, K_7 - 3P_2) = 18$ [SchSch2]

$$R(K_3, K_3 + \overline{K}_m) = R(K_3, K_3 + C_m) = 2m + 5 \text{ for } m \geq 212 \text{ [Zhou1]}$$

$$R(K_3, K_2 + T_n) = 2n + 3 \text{ for } n\text{-vertex trees } T_n, \text{ for } n \geq 4 \text{ [SoGQ]}$$

$R(K_3, G) = 2n(G) - 1$ for any connected G on at least 4 vertices and with at most $(17n(G) + 1)/15$ edges, in particular for $G = P_i$ and $G = C_i$, for all $i \geq 4$ [BEFRS1]

$R(K_3, G) \leq 2e(G) + 1$ for any graph G without isolated vertices [Sid3, GK]

$R(K_3, G) \leq n(G) + e(G)$ for all G , a conjecture [Sid2]

$R(K_3, G)$ for all connected G up to 9 vertices [BBH1, BBH2], see also section 7.1

$R(K_3, K_n)$, see section 2

$R(K_3, K_n - e)$, see section 3.1

Formulas for $R(nK_3, mG)$ for all G of order 4 without isolates [Zeng]

Since $B_1 = F_1 = C_3 = W_3 = K_3$, other sections apply

See also [AKS, BBH1, BBH2, FrLo, Fra1, Fra2, Gri, Loc, KM1, LiZa1, RK3, RK4, She2, Spe2, Stat, Yu1]

4.9. Cycles versus other graphs

Note: for cycles versus K_n, W_n, B_n - see section 3.

C_4 versus stars [Par3, Par4, Par5, BEFRS5, Chen, ChenJ, GoMC, MoCa]

C_4 versus trees [EFRS4, Bu7, BEFRS5, Chen]

C_4 versus all graphs on six vertices [JR3]

C_4 versus various types of complete bipartite graphs, see section 3.2

$R(C_4, G) \leq 2q + 1$ for any isolate-free graph G with q edges [RoJa2]

$R(C_4, G) \leq p + q - 1$ for any connected graph G on p vertices and q edges [RoJa2]

$R(C_5, K_6 - e) = 17$ [JR4]

$R(C_5, K_4 - e) = 9$ [CRSPS]

C_5 versus all graphs on six vertices [JR4]

$R(C_6, K_5 - e) = 17$ [JR2]

C_6 versus all graphs on five vertices [JR2]

$R(C_{2m+1}, G) = 2n - 1$ for sufficiently large sparse graphs G on n vertices, in particular

$R(C_{2m+1}, T_n) = 2n - 1$ for all $n > 1512m + 756$, for n vertex trees T_n [BEFRS2]

$R(C_n, G) \leq 2q + \lfloor n/2 \rfloor - 1$, for $3 \leq n \leq 5$, for any isolate-free graph G with $q > 3$ edges.

It is conjectured that it also holds for other n [RoJa2].

Cycles versus paths [FLPS, BEFRS2]

Cycles versus stars [La1, Clark, see Par6]

Cycles versus trees [BEFRS2, FSS1]

Monotone paths and cycles [Lef]

Cycles versus $K_{n,m}$ and multipartite complete graphs [BoEr]

4.10. Wheels versus other graphs

Notes: In this survey the wheel graph $W_n = K_1 + C_{n-1}$ has n vertices, while some authors use the definition $W_n = K_1 + C_n$ with $n + 1$ vertices. For cycles versus W_n - see section 3.4.

$$R(W_5, K_5 - e) = 17 \text{ [He2][YH]}$$

$$R(W_5, K_5) = 27 \text{ [He2][RST]}$$

$$R(W_5, K_6) \geq 33, R(W_5, K_7) \geq 43 \text{ [Shao]}$$

W_5 and W_6 versus stars and paths [SuBa1]

W_5 versus $nK_{1,m}$ [BaHA]

W_5 and W_6 versus trees [BSNM]

W_7 and W_8 versus paths [Bas]

W_7 versus trees T_n with $\Delta(T_n) \geq n - 3$, other special trees T , and for $n \leq 8$ [ChenZZ3, ChenZZ5, ChenZZ6]

W_7 and W_8 versus trees [ChenZZ4, ChenZZ5]

W_9 versus stars [Zhang2, ZhaCZ1, ChenCZ2]

W_9 versus trees of high degree [ZhaCZ2]

Wheels versus stars [HaBA1, ChenZZ2, Kor]

Wheels W_n , for even n , versus star-like trees [SuBB1]

Wheels versus paths [BaSu, ChenZZ1, SaBr3, Zhang1]

Wheels versus books [Zhou3]

Wheels versus unions of stars [BaHA, HaBA2]

Wheels versus linear forests (disjoint unions of paths) [SuBa2]

Upper bound asymptotics for $R(W_n, K_m)$ [Song5, SoBL]

4.11. Books versus other graphs

Note: for cycles versus B_n see section 3.5.

$$R(B_3, K_4) = 14 \text{ [He3]}$$

$$R(B_3, K_5) = 20 \text{ [He2][BaRT]}$$

$$R(B_4, K_{1,4}) = 11 \text{ [RS2]}$$

Cyclic lower bounds for $R(B_m, K_n)$ for $m \leq 7, n \leq 9$ and for $R(B_3, K_n - e)$ for $n \leq 7$ [Shao]

Books versus paths [RS2]

Books versus stars [CRSPS, RS2]

Books versus trees [EFRS7]

Books versus K_n [LR1, Sud2]

Books versus wheels [Zhou3]

Books versus $K_2 + C_n$ [Zhou3]

Books and $(K_1 + tree)$ versus K_n [LR1]

Generalized books $K_r + qK_1$ versus K_n [NiRo1, NiRo4]

4.12. Trees and forests versus other graphs

In this subsection T_n and F_n denote n -vertex tree and forest, respectively.

$$R(T_n, K_m) = (n-1)(m-1) + 1 \quad [\text{Chv}]$$

$$R(T_n, C_{2m+1}) = 2n - 1 \quad \text{for all } n > 1512m + 756 \quad [\text{BEFRS2}]$$

$$R(T_n, B_m) = 2n - 1 \quad \text{for all } n \geq 3m - 3 \quad [\text{EFRS7}]$$

$R(F_{nk}, K_m) = (n-1)(m-2) + nk$ for all forests F_{nk} consisting of k trees with n vertices each, also exact formula for all other cases of forests versus K_m [Stahl]

Exact results for almost all small ($n(G) \leq 5$) connected G versus all trees [FRS4]

Trees versus C_4 [EFRS4, Bu7, BEFRSS5, Chen]

Trees versus paths [FS4, FSS1]

Trees versus cycles [FSS1, EFRS6]

Trees versus stars [Bu1, Cheng, Coc, GuoV, ZZ1]

Trees versus books [EFRS7]

Trees versus W_5 and W_6 [BSNM]

Trees versus W_7 and W_8 [ChenZZ4, ChenZZ5]

Trees T_n with $\Delta(T_n) \geq n - 3$, other special trees T , and for $n \leq 8$ versus W_7 [ChenZZ3, ChenZZ5, ChenZZ6]

Trees T_n with $\Delta(T_n) \geq n - 4$ versus W_9 [ZhaCZ2]

Star-like trees versus odd wheels [SuBB1, ChenZZ3]

Trees versus $K_n + \bar{K}_m$ [RS2, FSR]

Trees versus bipartite graphs [BEFRS5, EFRS6]

Trees versus almost complete graphs [GoJa2]

Trees versus multipartite complete graphs [EFRS8, BEFRSGJ]

Linear forests versus wheels [SuBa2]

Forests versus almost complete graphs [CGP]

Forests versus complete graphs [BE1, Stahl, BaHA]

Study of graphs G for which all or almost all trees are G -good [BF, BEFRSGJ], see also section 4.17, item [Bu2], for the definition and more pointers.

See also various parts of this survey for special trees, sparse graphs and section 4.4.

4.13. Cases for $n(G), n(H) \leq 5$

Clancy [Clan], in 1977, presented a table of $R(G, H)$ for all isolate-free graphs G with $n(G) = 4$ and H with $n(H) = 4$, except 5 entries. All five of the open entries have been solved as follows:

$R(B_3, K_4) = 14$	[He3]
$R(K_4 - e, K_5) = 16$	[BH]
$R(W_5, K_4) = 17$	[He2]
$R(K_5 - e, K_4) = 19$	[EHM1]
$R(K_5, K_4) = R(4, 5) = 25$	[MR4]

An interesting case in [Clan] is

$$R(K_4, K_5 - P_3) = R(K_4, K_4 + e) = R(4, 4) = 18.$$

Hendry [He2], in 1989, presented a table of $R(G, H)$ for all graphs G and H on 5 vertices without isolates, except 7 entries. Four of the open entries have been solved:

$R(K_5, K_4 + e) = R(4, 5) = 25$	[Ka1][MR4]
$R(W_5, K_5 - e) = 17$	[He2][YH]
$R(B_3, K_5) = 20$	[He2][BaRT]
$R(W_5, K_5) = 27$	[He2][RST]

The still open cases for K_5 versus $K_5 - P_3$, $K_5 - e$ and K_5 are as follows:

$25 \leq R(K_5, K_5 - P_3) \leq 26$	[Ka1][BLR]
$30 \leq R(K_5, K_5 - e) \leq 34$	[Ex8][Ex8]
$43 \leq R(K_5, K_5) \leq 49$	[Ex4][MR5]

All critical colorings for the case $R(C_5 + e, K_5) = 17$ were found in [He5].

4.14. Mixed cases

- $26 \leq R(K_{2,2,2}, K_{2,2,2})$, $K_{2,2,2}$ is an octahedron [Ex8]
- Unicyclic graphs [Gros1, Köh, KrRod]
- $K_{2,m}$ and C_{2m} versus K_n [CLRZ]
- $K_{2,n}$ versus any graph [RoJa2]
- Union of two stars [Gros2]
- Double stars* [GHK]
- Graphs with bridge versus K_n [Li1]
- Multipartite complete graphs [BEFRS3, FRS3, Stev]
- Multipartite complete graphs versus sparse graphs [EFRS4]
- Multipartite complete graphs versus trees [EFRS8, BEFRSGJ]
- Graphs with long tails [Bu2, BG]
- Brooms+ [EFRS3]

* double star is a union of two stars with their centers joined by an edge

+ broom is a star with a path attached to its center

4.15. Multiple copies of graphs, disconnected graphs

$2K_2$ versus all isolate-free graphs [CH2]

nK_2 versus mK_2 , in particular $R(nK_2, nK_2) = 3n - 1$ for $n \geq 1$ [CL1, CL2, Lor]

nK_3 versus mK_3 , in particular $R(nK_3, nK_3) = 5n$ for $n \geq 2$ [BES], see also section 3.3

nK_3 versus mK_4 [LorMu]

$nK_{1,m}$ versus W_5 [BaHA]

$R(nK_4, nK_4) = 7n + 4$ for large n [Bu8]

Stripes mP_2 [CL1, CL2, Lor]

$R(mG, nH) \leq (m - 1)n(G) + (n - 1)n(H) + R(G, H)$ [BES]

Formulas for $R(nK_3, mG)$ for all isolate-free graphs G on 4 vertices [Zeng]

Variety of results for numbers $R(nG, mH)$ [Bu1, BES, HaBA2]

Disjoint unions of paths (also called linear forests) [BuRo2, FS2]

Disconnected graphs versus other graphs [BE1, GoJa1]

Study of Ramsey numbers for multiple copies of graphs [BES]

See section 3.3 for cases involving unions of cycles

See also [Bu9, BE1, LorMu, MiSa, Den, Biel1, Biel2]

4.16. General results for sparse graphs

[Chv] $R(K_n, T_m) = (n - 1)(m - 1) + 1$ for any tree T_m on m vertices.

[BEFRS2] $R(C_{2m+1}, G) = 2n - 1$ for sufficiently large sparse graphs G on n vertices, little more complicated formulas for P_{2m+1} instead of C_{2m+1} .

[BE3] Graphs yielding $R(K_n, G) = (n - 1)(n(G) - 1) + 1$, called Ramsey n -good, and related results (see also [EFRS5]). Recent extensive survey and further study of n -goodness appeared in [NiRo4].

[CRST] $R(G, G) \leq c_d n(G)$ for all G , where constant c_d depends only on the maximum degree d in G . The constant was improved in [GRR1, FoxSu1]. Tight lower and upper bounds for bipartite G [GRR2, Con2].

[BE1] Study of L -sets, which are sets of pairs of graphs whose Ramsey numbers are linear in the number of vertices. Conjecture that Ramsey numbers grow linearly for d -degenerate graphs (graph is d -degenerate if all its subgraphs have minimum degree at most d). Progress towards this conjecture was obtained by several authors, including [KoRö1, KoRö2, KoSu, FoxSu1, FoxSu2].

[ChenS] $R(G, G) \leq c_d n$ for all d -arrangeable graphs G on n vertices, in particular with the same constant for all planar graphs. The constant c_d was improved in [Eaton]. An extension to graphs not containing a subdivision of K_d [RöTh].

[Shi3] Ramsey numbers grow linearly for degenerate graphs versus some sparser graphs, arrangeable graphs, crowns, graphs with bounded maximum degree, planar graphs, and graphs without any topological minor of a fixed clique.

- [EFRS9] Study of graphs G , called *Ramsey size linear*, for which there exists a constant c_G such that for all H with no isolates $R(G, H) \leq c_G e(H)$. An overview and further results were given in [BaSS].
- [LRS] $R(G, G) < 6n$ for all n -vertex graphs G , in which no two vertices of degree at least 3 are adjacent. This improves the result $R(G, G) \leq 12n$ in [Alon1]. In an early paper [BE1] it was proved that if any two points of degree at least 3 are at distance at least 3 then $R(G, G) \leq 18n$.
- [Shi1] $R(Q_n, Q_n) \leq 2^{(3+\sqrt{5})n/2+o(n)}$, for the n -dimensional cube Q_n with 2^n vertices. This bound can also be derived from a theorem in [KoRö1]. An improvement was obtained in [Shi4], and a further one to $R(Q_n, Q_n) \leq 2^{2n+5n}$ in [FoxSu1].
- [Gros1] Conjecture that $R(G, G) = 2n(G) - 1$ if G is unicyclic of odd girth. Further support for the conjecture was given in [Köh, KrRod].
- [-] See also earlier subsections 4.* for various specific sparse graphs.

4.17. Other general two color results

- [CH2] $R(G, H) \geq (\chi(G) - 1)(c(H) - 1) + 1$, where $\chi(G)$ is the chromatic number of G , and $c(H)$ is the size of the largest connected component of H .
- [CH3] $R(G, G) > (s2^{e(G)-1})^{1/n(G)}$, where s is the number of automorphisms of G . Hence $R(K_{n,n}, K_{n,n}) > 2^n$, see also section 5.6.s.
- [BE2] $R(G, G) \geq \lfloor (4n(G) - 1)/3 \rfloor$ for any connected G , and $R(G, G) \geq 2n - 1$ for any connected nonbipartite G . These bounds can be achieved for all $n \geq 4$.
- [Bu2] Graphs H yielding $R(G, H) = (\chi(G) - 1)(n(H) - 1) + s(G)$, where $s(G)$ is a chromatic surplus of G , defined as the minimum number of vertices in some color class under all vertex colorings in $\chi(G)$ colors (such H 's are called G -good). This idea, initiated in [Bu2], is a basis of a number of exact results for $R(G, H)$ for large and sparse graphs H [BG, BEFRS2, BEFRS4, Bu5, FS, EFRS4, FRS3, BEFSRGJ, BF, LR4, Biel2]. Surveys of this area appeared in [FRS5, NiRo4].
- [BaLS] Graph G is Ramsey saturated if $R(G + e, G + e) > R(G, G)$ for every edge e in \bar{G} . Several theorems involving cycles, cycles with chords and trees on Ramsey saturated and unsaturated graphs. Seven conjectures including one stating that almost all graphs are Ramsey unsaturated. Special cases involving cycles and Jahangir graphs were studied in [AliSur].
- [Par3] Relations between some Ramsey graphs and block designs. See also [Par4].
- [Bra3] $R(G, H) > h(G, d)n(H)$ for all nonbipartite G and almost every d -regular H , for some h unbounded in d .
- [DoLL] Lower asymptotics of $R(G, H)$ depending on the average degree of G and the size of H . This continues the study initiated in [EFRS5], later much enhanced to both lower and upper bounds in [Sud3].

- [LiZa1] Lower bound asymptotics of $R(G, H)$ for large dense H .
- [AIKS] Discussion of a conjecture by Erdős that there exists a constant c such that $R(G, G) \leq 2^{c\sqrt{e(G)}}$. Proof for bipartite graphs G and progress towards the conjecture in other cases.
- [Kriv] Lower bound on $R(G, K_n)$ depending on the density of subgraphs of G . This construction for $G = K_m$ produces a bound similar to the best known probabilistic lower bound by Spencer [Spe2].
- [RoJa2] $R(K_{2,k}, G) \leq kq + 1$, for $k \geq 2$, for isolate-free graphs G with $q \geq 2$ edges.
- [FM] $R(W_6, W_6) = 17$ and $\chi(W_6) = 4$. This gives a counterexample $G = W_6$ to the Erdős conjecture (see [GRS]) $R(G, G) \geq R(K_{\chi(G)}, K_{\chi(G)})$, since $R(4, 4) = 18$.
- [BE1] $R(G + K_1, H) \leq R(K_{1, R(G, H)}, H)$.
- [LiShen] $R(\bar{K}_2 + G, \bar{K}_2 + G) \leq 4R(G, \bar{K}_2 + G) - 2$.
- [NiRo1] $R(K_{p+1}, B_q^r) = p(q+r-1) + 1$ for generalized books $B_q^r = K_r + qK_1$, for all sufficiently large q .
- [LR3] Bounds on $R(H + \bar{K}_n, K_n)$ for general H . Also, for fixed k and m , as $n \rightarrow \infty$, $R(K_k + \bar{K}_m, K_n) \leq (m + o(1))n^k / (\log n)^{k-1}$ [LiRZ1].
- [LiTZ] Asymptotics of $R(H + \bar{K}_n, K_n)$. In particular, the order of magnitude of $R(K_{m,n}, K_n)$ is $n^{m+1} / (\log n)^m$.
- [LiRZ2] Let G'' be a graph obtained from G by deleting two vertices. Then $R(G, H) \leq A + B + 2 + 2\sqrt{(A^2 + AB + B^2)/3}$, where $A = R(G'', H)$ and $B = R(G, H'')$.
- [BE1] Relations between the cases of G or $G + K_1$ versus H or $H + K_1$.
- [HaKr] Study of cyclic graphs yielding lower bounds for Ramsey numbers. Exact formulas for paths and cycles, small complete graphs and for graphs with up to five vertices.
- [Bu6] Given integer m and graphs G and H , determining whether $R(G, H) \leq m$ holds is NP-hard. Further complexity results related to Ramsey theory were presented in [Bu10].
- [Scha] Ramsey arrowing is Π_2^P -complete, a rare natural example of a problem higher than NP in the polynomial hierarchy of computational complexity theory.
- [-] Special cases of multicolor results listed in section 5.
- [-] See also surveys listed in section 7.

5. Multicolor Ramsey Numbers

The only known value of a multicolor classical Ramsey number:

$$R_3(3) = R(3,3,3) = R(3,3,3; 2) = 17 \quad \text{[GG]}$$

2 critical colorings (on 16 vertices) [KaSt, LayMa]

2 colorings on 15 vertices [Hein]

115 colorings on 14 vertices [PR1]

General upper bound, implicit in [GG]:

$$R(k_1, \dots, k_r) \leq 2 - r + \sum_{i=1}^r R(k_1, \dots, k_{i-1}, k_i - 1, k_{i+1}, \dots, k_r) \quad \text{(a)}$$

Inequality in (a) is strict if the right hand side is even, and at least one of the terms in the summation is even. It is suspected that this upper bound is never tight for $r \geq 3$ and $k_i \geq 3$, except for $r = k_1 = k_2 = k_3 = 3$. However, only two cases are known to improve over (a), namely $R_4(3) \leq 62$ [FKR] and $R(3,3,4) \leq 31$ [PR1, PR2], for which (a) produces only the bounds of 66 and 34, respectively.

5.1. Bounds for classical numbers

Diagonal Cases

<i>m</i>	3	4	5	6	7	8	9
<i>r</i>							
3	17 GG	128 HiIr	417 Ex17	1070 Mat	3214 XuR1	5384 XX2	13761 XXER
4	51 Chu1	634 XXER	3049 Xu	15202 XXER	62017 XXER		
5	162 Ex10	3416 XXER	26912 Xu				
6	538 FreSw						
7	1682 FreSw						

Table X. Known nontrivial lower bounds for diagonal multicolor Ramsey numbers $R_r(m)$, with references.

The best published bounds corresponding to the entries in Table X marked by personal communications [Ex17] and [Xu] are $415 \leq R_3(5)$, $2721 \leq R_4(5)$ and $26082 \leq R_5(5)$ [XXER].

The most studied and intriguing open case is

$$[\text{Chu1}] \quad 51 \leq R_4(3) = R(3,3,3,3) \leq 62 \quad [\text{FKR}]$$

The construction for $51 \leq R_4(3)$ as described in [Chu1] is correct, but be warned of a typo found by Christopher Frederick in 2003 (it makes a triangle (31,7,28) in color 1 in the displayed matrix). The inequality 5.a implies $R_4(3) \leq 66$, Folkman [Fo] in 1974 improved this bound to 65, and Sánchez-Flores [San] in 1995 proved $R_4(3) \leq 64$.

The upper bounds in $162 \leq R_5(3) \leq 307$, $538 \leq R_6(3) \leq 1838$, $1682 \leq R_7(3) \leq 12861$, $128 \leq R_3(4) \leq 236$ and $634 \leq R_4(4) \leq 6474$ are implied by 5.a (we repeat lower bounds from Table X just to easily see the ranges). All the latter and other upper bounds obtainable from known smaller bounds and 5.a can be computed with the help of a LISP program written by Kerber and Rowat [KerRo].

Off-Diagonal Cases

Three colors:

<i>m</i>	4	5	6	7	8	9	10	11	12	13	14
<i>k</i>											
3	30 Ka2	45 Ex2	60 Rob3	81 Ex16	101 Ex17	117 Ex17	141 5.2.c	157 5.2.c	182 LSS2	212 LSS2	233 5.2.c
4	55 KLR	81 Ex17	107 Ex17	143 Ex17	193 5.2.c						
5	81 Ex17	129 Ex17	169 5.2.c								

Table XI. Known nontrivial lower bounds for 3-color Ramsey numbers of the form $R(3, k, m)$, with references.

In addition, the bounds $303 \leq R(3,6,6)$, $609 \leq R(3,7,7)$ and $1689 \leq R(3,9,9)$ were derived in [XXER] (used there for building other lower bounds for some diagonal cases).

The other most studied, and perhaps the only open case of a classical multicolor Ramsey number, for which we can anticipate exact evaluation in the not-too-distance future is

$$[\text{Ka2}] \quad 30 \leq R(3,3,4) \leq 31 \quad [\text{PR1, PR2}]$$

In [PR1] it is conjectured that $R(3,3,4) = 30$, and the results in [PR2] eliminate some cases which could give $R(3,3,4) = 31$. The upper bounds in $45 \leq R(3,3,5) \leq 57$, $55 \leq R(3,4,4) \leq 79$, and $81 \leq R(3,4,5) \leq 160$ are implied by 5.(a) (we repeat lower bounds from the Table XI to show explicitly the current ranges).

Four colors:

$93 \leq R(3,3,3,4) \leq 153$	[Ex16, XXER], 5.a
$171 \leq R(3,3,4,4) \leq 462$	[Ex16, XXER], 5.a
$381 \leq R(3,4,4,4) \leq 1619$	5.2.g, 5.a
$162 \leq R(3,3,3,5)$	[XXER]
$561 \leq R(3,3,3,11)$	[XX2, XXER]

Lower bounds for higher numbers can be obtained by using general constructive results from section 5.2 below. For example, the bounds $261 \leq R(3,3,15)$ and $247 \leq R(3,3,3,7)$ were not published explicitly but are implied by 5.2.c and 5.2.d, respectively.

5.2. General results for complete graphs

- (b) $R_r(3) \geq 3R_{r-1}(3) + R_{r-3}(3) - 3$ [Chu1]
- (c) $R(3, k, l) \geq 4R(k, l-1) - 3$, and in general for $r \geq 2$ and $k_i \geq 2$
 $R(3, k_1, \dots, k_r) \geq 4R(k_1-1, k_2, \dots, k_r) - 3$ for $k_1 \geq 5$, and
 $R(k_1, 2k_2-1, k_3, \dots, k_r) \geq 4R(k_1-1, k_2, \dots, k_r) - 3$ for $k_1 \geq 5$ [XX2, XXER]
- (d) $R(3, 3, 3, k_1, \dots, k_r) \geq 3R(3, 3, k_1, \dots, k_r) + R(k_1, \dots, k_r) - 3$ [Rob2]
- (e) $R_k(3) \leq k!(e - e^{-1} + 3)/2 \approx 2.67k!$ [Wan], which improves the classical upper bound of $3k!$ [GRS]. For more work, mostly on asymptotics of $R_k(3)$, see [AbbH, Fre, Chu2, ChGri, GRS, GrRö, XXER]
- (f) $R(k_1, \dots, k_r) \geq S(k_1, \dots, k_r) + 2$, where $S(k_1, \dots, k_r)$ is the generalized Schur number [AbbH, Gi1, Gi2]. In particular, the special case $k_1 = \dots = k_r = 3$ has been widely studied [Fre, FreSw, Ex10, Rob3].
- (g) $R(k_1, \dots, k_r) \geq L(k_1, \dots, k_r) + 1$, where $L(k_1, \dots, k_r)$ is the maximal order of any cyclic (k_1, \dots, k_r) -coloring, which can be considered a special case of Schur partitions defining (symmetric) Schur numbers. Many lower bounds for Ramsey numbers were established by cyclic colorings. The following recurrence can be used to derive lower bounds for higher parameters. For $k_i \geq 3$
 $L(k_1, \dots, k_r, k_{r+1}) \geq (2k_{r+1} - 3)L(k_1, \dots, k_r) - k_{r+1} + 2$ [Gi2]
- (h) $R_r(m) \geq p + 1$ and $R_r(m+1) \geq r(p+1) + 1$ if there exists a K_m -free cyclotomic r -class association scheme of order p [Mat].
- (i) If the quadratic residues Paley graph Q_p of prime order $p = 4t + 1$ contains no K_k , then $R(s, k+1, k+1) \geq 4ps - 6p + 3$ [XXER].
- (j) $R_r(m) \geq c_m(2m-3)^r$, and some slight improvements of this bound for small values of m [AbbH, Gi1, Gi2, Song2].

- (k) $R_r(pq + 1) > (R_r(p + 1) - 1)(R_r(q + 1) - 1)$ [Abb1]
- (l) $R_r(pq + 1) > R_r(p + 1)(R_r(q + 1) - 1)$ for $p \geq q$ [XXER]
- (m) $R(p_1q_1 + 1, \dots, p_rq_r + 1) > (R(p_1 + 1, \dots, p_r + 1) - 1)(R(q_1 + 1, \dots, q_r + 1) - 1)$ [Song3]
- (n) $R_{r+s}(m) > (R_r(m) - 1)(R_s(m) - 1)$ [Song2]
- (o) $R(k_1, k_2, \dots, k_r) > (R(k_1, \dots, k_i) - 1)(R(k_{i+1}, \dots, k_r) - 1)$ in [Song1], see [XXER].
- (p) $R(k_1, k_2, \dots, k_r) > (k_1 + 1)(R(k_2 - k_1 + 1, k_3, \dots, k_r) - 1)$ [Rob4]
- (q) Further lower bound constructions, though with more complicated assumptions, were presented in [XX2, XXER].
- (r) Grolmusz [Grol1] generalized the classical constructive lower bound by Frankl and Wilson [FraWi] (section 2.3.r) to more colors and to hypergraphs [Grol3] (section 6).
- (s) Exact asymptotics of a very special but important case is known, namely $R(3, 3, n) = \Theta(n^3 \text{poly log } n)$ [AlRö]. For general upper bounds and more asymptotics see in particular [Chu4, ChGra2, ChGri, GRS, GrRö].

All lower bounds in (b) through (r) above are constructive. (d) generalizes (b), (m) generalizes both (k) and (o), and (o) generalizes (n). (l) is stronger than (k). Finally observe that the construction (m) with $q_1 = \dots = q_i = 1 = p_{i+1} = \dots = p_r$ is the same as (o).

5.3. Cycles

Three colors

- (a) The first larger paper in this area by Erdős, Faudree, Rousseau and Schelp [EFRS1] appeared in 1976. It gives several formulas and bounds for $R(C_m, C_n, C_k)$ and $R(C_m, C_n, C_k, C_l)$ for large m . For three colors [EFRS1] includes:

$$\begin{aligned}
 R(C_m, C_{2p+1}, C_{2q+1}) &= 4m - 3 \text{ for } p \geq 2, q \geq 1, \\
 R(C_m, C_{2p}, C_{2q+1}) &= 2(m + p) - 3 \text{ and} \\
 R(C_m, C_{2p}, C_{2q}) &= m + p + q - 2 \text{ for } p, q \geq 1 \text{ and large } m.
 \end{aligned}$$

- (b) Triple even cycles.

$$R_3(C_{2m}) \geq 4m \text{ for all } m \geq 2 \text{ [DzNS], see also (g) and (h) below.}$$

$R(C_n, C_n, C_n) = (2 + o(1))n$ for even n [FiŁu, GyRSS]. A similar more general result holds for slightly off-diagonal cases [FiŁu]. The diagonal case was improved to exactly $2n$ for large n [SiqSk].

- (c) Triple odd cycles.

$$\begin{aligned}
 R_3(C_{2m+1}) &= 8m + 1 \text{ for all sufficiently large } m, \text{ or equivalently} \\
 R(C_n, C_n, C_n) &= 4n - 3 \text{ for all sufficiently large odd } n \text{ [KoSS].}
 \end{aligned}$$

$R(C_n, C_n, C_n) \leq (4 + o(1))n$, with equality for odd n [Łuc]. It was conjectured by Bondy and Erdős, see [Erd2], that $R(C_n, C_n, C_n) \leq 4n - 3$ for $n \geq 4$. If true, then for all odd $n \geq 5$ we have $R(C_n, C_n, C_n) = 4n - 3$.

$m\ n\ k$	$R(C_m, C_n, C_k)$	references	general results
3 3 3	17	GG	page 29
3 3 4	17	ExRe	
3 3 5	21	Sun1+/Tse3	$5k - 4$ for $k \geq 5$, $m = n = 3$ [Sun1+]
3 3 6	26	Sun1+	
3 3 7	31	Sun1+	
3 4 4	12	Schu	
3 4 5	13	Sun1+/Rao/Tse3	
3 4 6	13	Sun1+/Tse3	
3 4 7	15	Sun1+/Tse3	
3 5 5	≥ 17	Tse3	
3 5 6	21	Sun1+	
3 5 7	25	Sun1+	
3 6 6			
3 6 7	21	Sun1+	
3 7 7			
4 4 4	11	BS	
4 4 5	12	Sun2+/Tse3	
4 4 6	12	Sun2+/Tse3	$k + 2$ for $k \geq 11$, $m = n = 4$ [Sun2+]
4 4 7	12	Sun2+/Tse3	values for $k = 8, 9, 10$ are 12, 13, 13 [Sun2+]
4 5 5	13	Tse3	
4 5 6	13	Sun1+	
4 5 7	15	Sun1+	
4 6 6	11	Tse3	
4 6 7	13	Sun1+/Tse3+	
4 7 7			
5 5 5	17	YR1	
5 5 6	21	Sun1+	
5 5 7	25	Sun1+	
5 6 6			
5 6 7	21	Sun1+	
5 7 7			
6 6 6	12	YR2	$R_3(C_{2q}) \geq 4q$ for $q \geq 2$ [DzNS]
6 6 7	15	Sun1+	see (a) for larger parameters
6 7 7			see (a) for larger parameters
7 7 7	25	FSS2	$R_3(C_{2q+1}) = 8q + 1$ for large m [KoSS]
8 8 8	16	Sun	

Table XII. Ramsey numbers $R(C_m, C_n, C_k)$ for $m, n, k \leq 7$ and $m = n = k = 8$.
 (Sun1+ abbreviates SunYWLX, Sun2+ abbreviates SunYLZ2,
 the work in [SunYWLX] and [SunYLZ2] is independent from [Tse3])

More colors

For results on $R_k(C_3) = R_k(K_3)$ see sections 5.1, 5.2.

$$\begin{array}{ll}
 R_4(C_4) = 18 & \text{[Ex2] [SunYLZ1]} \\
 18 \leq R_4(C_6) & \text{[SunYJLS]} \\
 27 \leq R_5(C_4) \leq 29 & \text{[LaWo1]} \\
 R_5(C_6) = 26 & \text{[SunYJLS] [SunYW]}
 \end{array}$$

$$21 \leq R(C_3, C_4, C_4, C_4) \leq 27 \quad [\text{XuR2}]$$

$$28 \leq R(C_3, C_3, C_4, C_4) \leq 36 \quad [\text{XuR2}]$$

$$49 \leq R(C_3, C_3, C_3, C_4) \quad 5.6.n$$

- (d) Formulas for $R(C_m, C_n, C_k, C_l)$ for large m [EFRS1].
- (e) $R_k(C_4) \leq k^2 + k + 1$ for all $k \geq 1$, $R_k(C_4) \geq k^2 - k + 2$ for all $k - 1$ which is a prime power [Ir, Chu2, ChGra1], and $R_k(C_4) \geq k^2 + 2$ for odd prime power k [LaWo1]. The latter was extended to any prime power k in [Ling, LaMu].
- (f) $R_k(C_{2m}) \geq (k + 1)m$ for odd k and $m \geq 2$, and $R_k(C_{2m}) \geq (k + 1)m - 1$ for even k and $m \geq 2$ [DzNS].
- (g) $R_k(C_{2m}) \geq 2(k - 1)(m - 1) + 2$ [SunYXL].
- (h) $R_k(C_{2m}) \geq k^2 + 2m - k$ for $2m \geq k + 1$ and prime power k [SunYJLS].
- (i) $R_k(C_{2m}) = \Theta(k^{m/m-1})$ for fixed $m = 2, 3$ and 5 [LiLih].
- (j) $R_k(C_{2m}) \leq 201km$ for $k \leq 10^m/201m$ [EG].
- (k) $R_k(C_5) \leq \sqrt{18^k k!}$ [Li3].
- (l) $2^k m < R_k(C_{2m+1}) \leq (k + 2)!(2m + 1)$ [BoEr].
A better upper bound $R_k(C_{2m+1}) < 2(k + 2)!m$ was obtained in [EG].
- (m) Asymptotic bounds for $R_k(C_n)$ [Bu1, GRS, ChGra2, LiLih].
- (n) Survey of multicolor cycle cases [Li2].

Cycles versus other graphs

The bounds for the following six cases were established in [XSR1]:

$$19 \leq R(C_4, C_4, K_4) \leq 22 \quad 31 \leq R(C_4, C_4, C_4, K_4) \leq 50$$

$$25 \leq R(C_3, C_4, K_4) \leq 32 \quad 42 \leq R(C_3, C_4, C_4, K_4) \leq 76$$

$$52 \leq R(C_4, K_4, K_4) \leq 72 \quad 87 \leq R(C_4, C_4, K_4, K_4) \leq 179$$

$$R(K_{1,3}, C_4, K_4) = 16 \quad [\text{KM2}]$$

$$R(C_4, C_4, C_4, T) = 16 \text{ for } T = P_4 \text{ and } T = K_{1,3} \quad [\text{ExRe}]$$

- (o) Study of $R(C_n, K_{t_1}, \dots, K_{t_k})$ and $R(C_n, K_{t_1, s_1}, \dots, K_{t_k, s_k})$ for large n [EFRS1].
- (p) Study of asymptotics for $R(C_m, \dots, C_m, K_n)$, in particular for fixed number of colors we have $R(C_4, C_4, \dots, C_4, K_n) = \Theta(n^2/\log^2 n)$ [AIRö].
- (q) Study of asymptotics for $R(C_{2m}, C_{2m}, K_n)$ for fixed m [AIRö, ShiuLL], in particular $R(C_4, C_4, K_n) = \Theta(n^2 \text{ poly log } n)$ [AIRö].
- (r) Monotone paths and cycles [Lef].
- (s) For combinations of C_3 and K_n see sections 2, 5.1 and 5.2.

5.4. Paths, paths versus other graphs

In 2007, Gyárfás, Ruszinkó, Sárközy and Szemerédi [GyRSS] established that for all sufficiently large n we have

$$R(P_n, P_n, P_n) = 2n - 2 + n \bmod 2.$$

Three color path and path-cycle cases

- (a) $R(P_m, P_n, P_k) = m + \lfloor n/2 \rfloor + \lfloor k/2 \rfloor - 2$ for $m \geq 6(n+k)^2$ [FS2],
It holds asymptotically for $m \geq n \geq k$ with an extra term $o(m)$ [FiŁu].
 $R_3(P_4) = 6$ [Ir], for other combinations of $3 \leq m, n, k \leq 4$ the value is 5 [AKM].
- (b) $R(P_3, P_3, C_m) = 5, 6, 6$, for $m = 3, 4$ [AKM], 5, and
 $R(P_3, P_3, C_m) = m$ for $m \geq 6$ [Dzi].
 $R(P_3, P_4, C_m) = 7$ for $m = 3, 4$ [AKM] and 5, and
 $R(P_3, P_4, C_m) = m + 1$ for $m \geq 6$ [Dzi].
 $R(P_4, P_4, C_m) = 9, 7, 9$ for $m = 3, 4$ [AKM] and 5 [Dzi], and
 $R(P_4, P_4, C_m) = m + 2$ for $m \geq 6$ [DzKP].
- (c) $R(P_3, P_5, C_m) = 9, 7, 9, 7, 9$ for $m = 3, 4, 5, 6, 7$ [Dzi, DzFi2], and
 $R(P_3, P_5, C_m) = m + 1$ for $m \geq 8$ [DzKP].
A table of $R(P_3, P_k, C_m)$ for all $3 \leq k \leq 8$ and $3 \leq m \leq 9$ [DzFi2].
- (d) $R(P_4, P_5, C_m) = 11, 7, 11, 11$, and $m + 2$ for $m = 3, 4, 5, 7$ and $m \geq 23$, and
 $R(P_4, P_6, C_m) = 13, 8, 13, 13$, and $m + 3$ for $m = 3, 4, 5, 7$ and $m \geq 18$ [ShaXSP].
- (e) $R(P_3, P_n, C_4) = n + 1$ for $n \geq 6$,
 $R(P_m, P_n, C_k) = 2n + 2\lfloor m/2 \rfloor - 3$ for large n and odd $m \geq 3$, in particular
 $R(P_3, P_n, C_k) = 2n - 1$,
 $R(P_4, P_n, C_k) = 2n + 1$ for odd $m \geq 3$ and $k \geq m$ [DzFi2].
- (f) $R(P_3, C_3, C_3) = 11$ [BE3], $R(P_3, C_4, C_4) = 8$ [AKM], $R(P_3, C_6, C_6) = 9$ [Dzi],
 $R(P_3, C_m, C_m) = R(C_m, C_m) = 2m - 1$ for odd $m \geq 5$ [DzKP] (for $m = 5, 7$ [Dzi]).
 $R(P_3, C_3, C_4) = 8$ [AKM], $R(P_3, C_3, C_5) = 9$, $R(P_3, C_3, C_6) = 11$,
 $R(P_3, C_3, C_7) = 13$, $R(P_3, C_4, C_5) = 8$, $R(P_3, C_4, C_6) = 8$,
 $R(P_3, C_4, C_7) = 8$, $R(P_3, C_5, C_6) = 11$, $R(P_3, C_5, C_7) = 13$ and
 $R(P_3, C_6, C_7) = 11$ [Dzi].
- (g) Formulas for $R(pP_3, qP_3, rP_3)$ and $R(pP_4, qP_4, rP_4)$ [Scob].
- (h) $R(P_3, K_4 - e, K_4 - e) = 11$ [Ex7]. All colorings (which can be any color neighborhood for the open case $R_3(K_4 - e)$, see section 5.5) were found in [Piw2].

More colors

- (i) $R_k(P_3) = k + 1 + (k \bmod 2)$, $R_k(2P_2) = k + 3$ for all $k \geq 1$ [Ir].
- (j) $R_k(P_4) = 2k + c_k$ for all k and some $0 \leq c_k \leq 2$. If k is not divisible by 3 then $c_k = 3 - k \bmod 3$ [Ir]. Wallis [Wall] showed $R_6(P_4) = 13$, which already implied $R_{3t}(P_4) = 6t + 1$, for all $t \geq 2$. Independently, the case $R_k(P_4)$ for $k \neq 3^m$ was completed by Lindström in [Lind], and later Bierbrauer proved $R_{3^m}(P_4) = 2 \cdot 3^m + 1$ for all $m > 1$. $R_3(P_4) = 6$ [Ir].
- (k) Formula for $R(P_{n_1}, \dots, P_{n_k})$ for large n_1 [FS2].
- (l) Formula for $R(n_1P_2, \dots, n_kP_2)$, in particular $R(nP_2, nP_2, nP_2) = 4n - 2$ [CL1].
- (m) Cockayne and Lorimer [CL1] found the exact formula for $R(n_1P_2, \dots, n_kP_2)$, and later Lorimer [Lor] extended it to a more general case of $R(K_m, n_1P_2, \dots, n_kP_2)$. More general cases of the latter, with multiple copies of the complete graph, stars and forests, were studied in [Stahl, LorSe, LorSo, GyRSS].
- (n) Multicolor cases for one large path or cycle involving small paths, cycles, complete and complete bipartite graphs [EFRS1].
- (o) See section 7.2, especially [AKM], for a number of cases for triples of small graphs.

5.5. Special cases

$$\begin{array}{ll}
 R_3(K_3 + e) = R_3(K_3) & [=17] \quad \text{[YR3, AKM]} \\
 28 \leq R_3(K_4 - e) \leq 30 & \text{[Ex7] [Piw2]} \\
 R(K_4 - e, K_4 - e, P_3) = 11 & \text{[Ex7]}
 \end{array}$$

All colorings for $(K_4 - e, K_4 - e, P_3)$ were found in [Piw2].

5.6. Other general results

- (a) Formulas for $R_k(G)$, where G is one of the graphs P_3 , $2K_2$ and $K_{1,3}$ for all k , and for P_4 if k is not divisible by 3 [Ir]. For some details see section 5.4.j.
- (b) $tk^2 + 1 \leq R_k(K_{2,t+1}) \leq tk^2 + k + 2$, where the upper bound is general, and the lower bound holds when both t and k are prime powers [ChGra1, LaMu].
- (c) $(m-1)\lfloor (k+1)/2 \rfloor < R_k(T_m) \leq 2km + 1$ for any tree T_m with m edges [EG], see also [GRS]. The lower bound can be improved for special large k [EG, GRS]. The upper bound was improved to $R_k(T_m) < (m-1)(k + \sqrt{k(k-1)}) + 2$ in [GyTu].
- (d) $k(\sqrt{m} - 1)/2 < R_k(F_m) < 4km$ for any forest F_m with m edges [EG], see [GRS]. See also pointers in items (u) and (v) below.
- (e) $R(m_1G_1, \dots, m_kG_k) \leq R(G_1, \dots, G_k) + \sum_{i=1}^k n(G_i)(m_i - 1)$, exercise 8.3.28 in [West].

- (f) Formula for $R(S_1, \dots, S_k)$, where S_i 's are arbitrary stars [BuRo1], see also [GauST].
- (g) Formula for $R(S_1, \dots, S_k, K_n)$, where S_i 's are arbitrary stars [Jac].
- (h) Formula for $R(S_1, \dots, S_k, nK_2)$, where S_i 's are arbitrary stars [CL2].
- (i) Formula for $R(S_1, \dots, S_k, T)$, where S_i 's are stars and T is a tree [ZZ1].
- (j) Formulas for $R(S_1, \dots, S_k)$, where each S_i 's is a star or $m_i K_2$ [ZZ2, EG].
- (k) Study of the case $R(K_m, n_1 P_2, \dots, n_k P_2)$ [Lor]. More general cases, with multiple copies of the complete graph, stars and forests, were investigated in [Stahl, LorSe, LorSo, GyRSS]. See also section 5.4.
- (l) If G is connected and $R(K_k, G) = (k-1)(n(G)-1) + 1$, in particular if G is any tree, then $R(K_{k_1}, \dots, K_{k_r}, G) = (R(k_1, \dots, k_r) - 1)(n(G) - 1) + 1$ [BE3]. A generalization for connected G_1, \dots, G_n in place of G appeared in [Jac].
- (m) If F, G, H are connected graphs then $R(F, G, H) \geq (R(F, G) - 1)(\chi(H) - 1) + \min\{R(F, G), s(H)\}$, where $s(G)$ is the chromatic surplus of G (see item [Bu2] in section 4.16). This leads to several formulas and bounds for F and G being stars and/or trees when $H = K_n$ [ShiuLL].
- (n) $R(K_{k_1}, \dots, K_{k_r}, G_1, \dots, G_s) \geq (R(k_1, \dots, k_r) - 1)(R(G_1, \dots, G_s) - 1) + 1$ for arbitrary graphs G_1, \dots, G_s [Bev]. This generalizes 5.2.o.
- (o) Constructive bound $R(G_1, \dots, G_{t^{n-1}}) \geq t^n + 1$ for decompositions of K_{t^n} [LaWo1, LaWo2].
- (p) $R(G_1, \dots, G_k) \leq 32\Delta k^\Delta n$, where $n \geq n(G_i)$ and $\Delta \geq \Delta(G_i)$ for all $1 \leq i \leq k$ [FoxSu1].
- (q) $R(G_1, \dots, G_k) \leq k^{2k\Delta q} n$, where $q \geq \chi(G_i)$ for all $1 \leq i \leq k$ [FoxSu1].
- (r) Bounds on $R_k(G)$ for unicyclic graphs G of odd girth.
Some exact values for special graphs G , for $k = 3$ and $k = 4$ [KrRod].
- (s) $R_k(G) > (sk^{e(G)-1})^{1/n(G)}$, where s is the number of automorphisms of G [CH3]. Other general bounds for $R_k(G)$ [CH3, Par6].
- (t) Bounds on $R_k(K_{s,t})$, in particular for $K_{2,2} = C_4$ and $K_{2,t}$ [ChGra1, AFM]. Asymptotics of $R_k(K_{s,t})$ for fixed k and s [LiTZ]. Upper bounds on $R_k(K_{s,t})$ [SunLi].
- (u) Bounds on $R_k(G)$ for trees, forests, stars and cycles [Bu1].
- (v) Bounds for trees $R_k(T)$ and forests $R_k(F)$ [EG, GRS, BB, GyTu, Bra1, Bra2, SwPr].
- (x) Study of $R(G_1, \dots, G_k, G)$ for large sparse G [EFRS1, Bu3].
- (y) Study of asymptotics for $R(C_n, \dots, C_n, K_m)$ [AlRö]. See also sections 5.3.pq.
- (z) See section 7.2, especially [AKM], for a number of cases for other small graphs, similar to those listed in section 5.3. See also surveys listed in section 7.

6. Hypergraph Numbers

The only known value of a classical Ramsey number for hypergraphs:

$$R(4,4;3) = 13$$

more than 200000 critical colorings [MR1]

The computer evaluation of $R(4,4;3)$ in 1991 consisted of an improvement of the upper bound from 15 to 13. This result followed an extensive theoretical study of this number by several authors [Gi4, Isb1, Sid1].

Special hypergraph cases

$$33 \leq R(4,5;3) \quad [\text{Ex13}]$$

$$38 \leq R(4,6;3) \quad [\text{HuSo+}]$$

$$65 \leq R(5,5;3) \quad [\text{Ea1}]$$

$$56 \leq R(4,4,4;3) \quad [\text{Ex8}]$$

$$34 \leq R(5,5;4) \quad [\text{Ex11}]$$

$$R(K_4-t, K_4-t;3) = 7 \quad [\text{Ea2}]$$

$$R(K_4-t, K_4;3) = 8 \quad [\text{Sob, Ex1, MR1}]$$

$$14 \leq R(K_4-t, K_5;3) \quad [\text{Ex1}]$$

$$13 \leq R(K_4-t, K_4-t, K_4-t;3) \leq 17 \quad [\text{Ex1}][\text{Ea1}]$$

- (a) The first bound on $R(4,5;3) \geq 24$ was obtained by Isbell [Isb2]. Shastri [Shas] gave a weak bound $R(5,5;4) \geq 19$ (now 34 in [Ex11]), nevertheless his lemmas, the stepping-up lemmas by Erdős and Hajnal (see [GRS, GrRö], also 6.j below), and others in [Ka3, Abb2, GRS, GrRö, HuSo, SoYL] can be used to derive better lower bounds for higher numbers.
- (b) Several lower bound constructions for 3-uniform hypergraphs were presented in [HuSo]. Study of lower bounds on $R(p,q;4)$ can be found in [Song3] and [SoYL, Song4] (the latter two papers are almost the same in contents). Most lower bounds in these papers can be easily improved by using the same techniques, but starting with better constructions for small parameters as listed above.
- (c) $R(p,q;4) \geq 2R(p-1,q;4) - 1$ for $p,q > 4$, and
 $R(p,q;4) \geq (p-1)R(p-1,q;4) - p + 2$ for $p \geq 5, q \geq 7$ [SoYL].

General results for 3-uniform hypergraphs

- (d) $2^{cn^2} < R(n,n;3) < 2^{2^n}$, Erdős, Hajnal and Rado (see [ChGra2] p. 30).
- (e) For some a,b the numbers $R(m,a,b;3)$ are at least exponential in m [AbbS].
- (f) $R(G,G;3) \leq c \cdot n(H)$ for some constant c depending only on the maximum degree of a 3-uniform hypergraph H [CooFKO, NaORS]. Similar results were proved for r -uniform hypergraphs in [KüCFO, Ishi, ConFS3], see (n) below.

- (g) A *loose* 3-uniform cycle C_n on $[n]$ is the set of triples $\{123, 345, 567, \dots, (n-1)n1\}$ (note that n must be even). For such loose cycles we have $R(C_{4k}, C_{4k}; 3) > 5k - 2$ and $R(C_{4k+2}, C_{4k+2}; 3) > 5k + 1$, and asymptotically these lower bounds are tight [HaŁP1+]. Generalizations to r -uniform hypergraphs and graphs other than cycles [GySS].
- (h) Study of the cases for 3-uniform *tight* cycles and paths in which consecutive edges share two points. For tight cycles, $R(C_{3k}, C_{3k}; 3) \approx 4k$ and $R(C_{3k+i}, C_{3k+i}; 3) \approx 6k$ for $i = 1$ or 2 , and for tight paths $R(P_k, P_k; 3) \approx 4k/3$ [HaŁP2+]. Related results in [PoRRS].
- (i) Lower and upper asymptotics of $R(s, n; k)$ for fixed s , and lower bound on $R_3(n; 3)$ [ConFS1]. Upper bounds on $R_k(H; 3)$ for complete multipartite 3-uniform hypergraphs H , a 4-color case, and some other general and special cases [ConFS2, ConFS3].

General results for hypergraphs

- (j) If $R(n, n; k) > m$ then $R(2n + k - 4, 2n + k - 4; k + 1) > 2^m$ for $n > k \geq 3$ (see [GRS] p. 106). This is the so-called stepping-up lemma, usually credited to Erdős and Hajnal.
- (k) Lower bounds on $R_k(n; r)$ are discussed in [AbbW, DLR].
- (l) General lower bounds for large number of colors were given in an early paper by Hirschfeld [Hir], and some of them were later improved in [AbbL].
- (m) Exact results for large 2-loose cycles (generalizing (g) above) and 2- and 3-color cases for all r -uniform diamond matchings [GySS].
- (n) $R(H, H; r) \leq c \cdot n(H)^{1+\varepsilon}$, for some constant $c = c(\Delta, r, \varepsilon)$ depending only on the maximum degree of H , r and $\varepsilon > 0$ [KoRö3]. The proofs of the linear bound $c \cdot n(H)$ were obtained independently in [KüCFO] and [Ishi], the latter including the multicolor case, and then without regularity lemma in [ConFS3]. More discussion of lower and upper bounds for various cases can be found in [ConFS1, ConFS2, ConFS3].
- (o) Let $H^{(r)}(s, t)$ be the complete r -partite r -uniform hypergraph with $r - 2$ parts of size 1, one part of size s , and one part of size t (for example, for $r = 2$ it is the same as $K_{s, t}$). For the multicolor numbers, Lazebnik and Mubayi [LaMu] proved that

$$tk^2 - k + 1 \leq R_k(H^{(r)}(2, t+1); r) \leq tk^2 + k + r,$$

where the lower bound holds when both t and k are prime powers. For the general case of $H^{(r)}(s, t)$, more bounds are presented in [LaMu].

- (p) Grolmusz [Grol1] generalized the classical constructive lower bound by Frankl and Wilson [FraWi] (section 2.3.s) to more colors and to hypergraphs [Grol3].
- (q) Lower and upper asymptotics, and other theoretical results on hypergraph numbers are gathered in [GrRö, GRS, ConFS1, ConFS2, ConFS3].

7. Cumulative Data and Surveys

7.1. Cumulative data for two colors

- [CH1] $R(G, G)$ for all graphs G without isolates on at most 4 vertices.
- [CH2] $R(G, H)$ for all graphs G and H without isolates on at most 4 vertices.
- [Clan] $R(G, H)$ for all graphs G on at most 4 vertices and H on 5 vertices, except five entries (now all solved, see section 4.13). All critical colorings for the isolate-free graphs G and H studied in [Clan] were found in [He4].
- [Bu4] $R(G, G)$ for all graphs G without isolates and with at most 6 edges.
- [He1] $R(G, G)$ for all graphs G without isolates and with at most 7 edges.
- [HaMe2] $R(G, G)$ for all graphs G on 5 vertices and with 7 or 8 edges.
- [He2] $R(G, H)$ for all graphs G and H on 5 vertices without isolates, except 7 entries (3 still open, see 4.13 and the paragraph at the end of this section).
- [HoMe] $R(G, H)$ for $G = K_{1,3} + e$ and $G = K_4 - e$ versus all connected graphs H on 6 vertices, except $R(K_4 - e, K_6)$. The result $R(K_4 - e, K_6) = 21$ was claimed by McNamara [McN, unpublished].
- [FRS4] $R(G, T)$ for all connected graphs G with $n(G) \leq 5$, and almost all trees T .
- [FRS1] $R(K_3, G)$ for all connected graphs G on 6 vertices.
- [Jin] $R(K_3, G)$ for all connected graphs G on 7 vertices.
Some errors in [Jin] were found [SchSch1].
- [Zeng] Formulas for $R(nK_3, mG)$ for all G of order 4 without isolates.
- [Brin] $R(K_3, G)$ for all connected graphs G on at most 8 vertices. The numbers for K_3 versus sets of graphs with fixed number of edges, on at most 8 vertices, were presented in [KM1].
- [BBH1] $R(K_3, G)$ for all connected graphs G on 9 vertices. See also [BBH2].
- [JR3] $R(C_4, G)$ for all graphs G on at most 6 vertices.
- [JR4] $R(C_5, G)$ for all graphs G on at most 6 vertices.
- [JR2] $R(C_6, G)$ for all graphs G on at most 5 vertices.
- [LoM3] $R(K_{2,n}, K_{2,m})$ for all $2 \leq n, m \leq 10$ except 8 cases, for which lower and upper bounds are given. Further data for other complete bipartite graphs in section 3.2 and [LoMe4].
- [HaKr] All best lower bounds up to 102 from cyclic graphs. Formulas for best cyclic lower bounds for paths and cycles, small complete graphs and for graphs with up to five vertices.

Chvátal and Harary [CH1, CH2] formulated several simple but very useful observations how to discover values of some numbers. All five missing entries in the tables of Clancy [Clan] have been solved (section 4.13). Out of 7 open cases in [He2] 4 have been solved, including

$R(4, 5) = R(G_{19}, G_{23}) = 25$ and other items listed in section 4.13. The still open 3 cases are for K_5 versus the graphs K_5 (section 2.1), $K_5 - e$ (section 3.1), and $K_5 - P_3$ (section 4.13).

7.2. Cumulative data for three colors

- [YR3] $R_3(G)$ for all graphs G with at most 4 edges and no isolates.
- [YR1] $R_3(G)$ for all graphs G with 5 edges and no isolates, except $K_4 - e$.
The case of $R_3(K_4 - e)$ remains open (see section 5.3).
- [YY] $R_3(G)$ for all graphs G with 6 edges and no isolates, except 10 cases.
- [AKM] $R(F, G, H)$ for most triples of isolate-free graphs with at most 4 vertices.
Some of the missing cases completed in [KM2].
- [DzFi2] $R(P_3, P_k, C_m)$ for all $3 \leq k \leq 8$ and $3 \leq m \leq 9$.

7.3. Surveys

- [Bu1] A general survey of results in Ramsey graph theory by S. A. Burr (1974)
- [Par6] A general survey of results in Ramsey graph theory by T. D. Parsons (1978)
- [BuRo3] Survey of results and new problems on multiplicities and Ramsey multiplicities by S. A. Burr and V. Rosta (1980)
- [Har2] Summary of progress by Frank Harary (1981)
- [ChGri] A general survey of bounds and values by F. R. K. Chung and C. M. Grinstead (1983)
- [JGT] Special volume of the *Journal of Graph Theory* (1983)
- [Rob1] A review of Ramsey graph theory for newcomers by F. S. Roberts (1984)
- [Bu7] What can we hope to accomplish in generalized Ramsey Theory? (1987)
- [GrRö] Survey of asymptotic problems by R. L. Graham and V. Rödl (1987)
- [GRS] An excellent book by R. L. Graham, B. L. Rothschild and J. H. Spencer, second edition (1990)
- [FRS5] Survey by Faudree, Rousseau and Schelp of graph goodness results, i.e. conditions for the formula $R(G, H) = (\chi(G) - 1)(n(H) - 1) + s(G)$ (1991)
- [Neš] A chapter in *Handbook of Combinatorics* by J. Nešetřil (1996)
- [Caro] Survey of zero-sum Ramsey theory by Y. Caro (1996)
- [Chu4] Among 114 open problems and conjectures of Paul Erdős, presented and commented by F. R. K. Chung, 31 are concerned directly with Ramsey numbers. 216 references are given (1997). An extended version of this work was prepared jointly with R. L. Graham [ChGra2]. (1998)
- [West] An extensive chapter on Ramsey theory in the very popular student textbook and researcher's manual of graph theory (2001)

- [GrNe] Ramsey Theory and Paul Erdős (2002)
- [CoPC] Special issue of *Combinatorics, Probability and Computing* (2003)
- [Ros2] Dynamic survey of Ramsey theory applications by V. Rosta (2004). A website maintained by W. Gasarch [Gas] gathers over 50 pointers to literature on applications of Ramsey theory in computer science. (2009)
- [Soi] History, results and people of Ramsey theory. The mathematical coloring book, mathematics of coloring and the colorful life of its creators. (2009)

The surveys by S. A. Burr [Bu1] and T. D. Parsons [Par6] contain extensive chapters on general exact results in graph Ramsey theory. F. Harary presented the state of the theory in 1981 in [Har2], where he also gathered many references including seven to other early surveys of this area. More than two decades ago, Chung and Grinstead in their survey paper [ChGri] gave less data than in this work, but included a broad discussion of different methods used in Ramsey computations in the classical case. S. A. Burr, one of the most experienced researchers in Ramsey graph theory, formulated in [Bu7] seven conjectures on Ramsey numbers for sufficiently large and sparse graphs, and reviewed the evidence for them found in the literature. Three of them have been refuted in [Bra3].

For newer extensive presentations see [GRS, GrRö, FRS5, Neš, Chu4, ChGra2], though these focus on asymptotic theory not on the numbers themselves. A very welcome addition is the 2004 compilation of applications of Ramsey theory by Rosta [Ros2]. This survey could not be complete without recommending special volumes of the *Journal of Graph Theory* [JGT, 1983] and *Combinatorics, Probability and Computing* [CoPC, 2003], which, besides a number of research papers, include historical notes and present to us Frank P. Ramsey (1903-1930) as a person. Finally, read the newest book by A. Soifer [Soi, 2009] on history and results in Ramsey theory!

8. Concluding Remarks

This compilation does not include information on numerous variations of Ramsey numbers, nor related topics, like size Ramsey numbers, zero-sum Ramsey numbers, irredundant Ramsey numbers, induced Ramsey numbers, local Ramsey numbers, connected Ramsey numbers, chromatic Ramsey numbers, avoiding sets of graphs in some colors, coloring graphs other than complete, or the so called Ramsey multiplicities. Interested readers can find such information in the surveys listed in section 7 here.

Ramsey@Home [RaHo] is a distributed computing project at the University of Wisconsin-Oshkosh designed to find new lower bounds for various Ramsey numbers. Join and help! Readers may be interested in knowing that the US patent 6965854 B2 issued on November 15, 2005 claims a method of using Ramsey numbers in "Methods, Systems and Computer Program Products for Screening Simulated Traffic for Randomness". Check the original document at <http://www.uspto.gov/patft> if you wish to find out whether your usage of Ramsey numbers is covered by this patent.

Acknowledgement

In addition to the many individuals who helped to improve consecutive versions of this survey, the author would like to specially thank Brendan McKay, Geoffrey Exoo and Heiko Harborth for their help in gathering data for the first versions. Thanks to the many others who over the years have helped me in the development and improvement of new revisions.

The author apologizes for any omissions or other errors in reporting results belonging to the scope of this work. Suggestions for any kind of corrections or additions will be greatly appreciated and considered for inclusion in the next revision of this survey.

References

Papers containing results obtained with the help of computer algorithms have been marked with stars. We identify two such categories of papers: those marked with * involving some use of computers where the results are easily verifiable with some computations, and those marked with ** where cpu intensive algorithms have to be implemented to replicate or verify the results. The first category contains mostly constructions done by algorithms, while the second mostly nonexistence results or claims of complete enumerations of special classes of graphs.

The references are ordered alphabetically by the last name of the first author, and where multiple papers have the same first author they are ordered by the last name of the second author, etc. We preferred that all work by the same author be in consecutive positions. Unfortunately, this causes that some of the abbreviations are not in alphabetical order. For example, [BaRT] is earlier on the list than [BaLS]. We also wish to explain a possible confusion with respect to the order of parts and spelling of Chinese names. We put them without any abbreviations, in most cases with the last name written first as is customary in original. This is often distinct from citations in other sources. One can obtain all variations of writing any specific name by consulting the authors database of *Mathematical Reviews* at <http://www.ams.org/mathscinet/search>.

A, B, C	pages 44 - 49
D, E, F	pages 49 - 53
G, H, I, J	pages 53 - 57
K, L	pages 57 - 61
M, N, O, P, Q, R	pages 61 - 64
S	pages 65 - 69
T, U, V, W, X, Y, Z	pages 69 - 72

A

- [Abb1] H.L. Abbott, *Ph. D. thesis*, University of Alberta, Edmonton, 1965.
- [Abb2] H.L. Abbott, A Theorem Concerning Higher Ramsey Numbers, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds.) Vol. 1, 25-28, Colloq. Math. Soc. Janos Bolyai, Vol. 10, North-Holland, Amsterdam, 1975.
- [AbbH] H.L. Abbott and D. Hanson, A Problem of Schur and Its Generalizations, *Acta Arithmetica*, **20** (1972) 175-187.
- [AbbL] H.L. Abbott and Andy Liu, Remarks on a Paper of Hirschfeld Concerning Ramsey Numbers, *Discrete Mathematics*, **39** (1982) 327-328.
- [AbbS] H.L. Abbott and M.J. Smuga-Otto, Lower Bounds for Hypergraph Ramsey Numbers, *Discrete Applied Mathematics*, **61** (1995) 177-180.
- [AbbW] H.L. Abbott and E.R. Williams, Lower Bounds for Some Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **16** (1974) 12-17.
- [AKS] M. Ajtai, J. Komlós and E. Szemerédi, A Note on Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **29** (1980) 354-360.
- [AliBas] K. Ali and E.T. Baskoro, On the Ramsey Numbers for a Combination of Paths and Jahangirs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **65** (2008) 113-119.
- [AliBT] K. Ali, E.T. Baskoro and I. Tomescu, On the Ramsey Numbers for Paths and Generalized Jahangir Graphs $J_{s,m}$, *Bull. Math. Soc. Sci. Math. R. S. Roumanie (N.S.)*, **51(99)** (2008) 177-182.
- [AliSur] K. Ali and Surahmat, A Cycle or Jahangir Ramsey Unsaturated Graphs, *Journal of Prime Research in Mathematics*, **2** (2006) 187-193.
- [Alon1] N. Alon, Subdivided Graphs Have Linear Ramsey Numbers, *Journal of Graph Theory*, **18** (1994) 343-347.
- [Alon2] N. Alon, Explicit Ramsey Graphs and Orthonormal Labelings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R12, **1** (1994), 8 pages.
- [AlKS] N. Alon, M. Krivelevich and B. Sudakov, Turán Numbers of Bipartite Graphs and Related Ramsey-Type Questions, *Combinatorics, Probability and Computing*, **12** (2003) 477-494.
- [AlPu] N. Alon and P. Pudlák, Constructive Lower Bounds for off-diagonal Ramsey Numbers, *Israel Journal of Mathematics*, **122** (2001) 243-251.
- [AlRö] N. Alon and V. Rödl, Sharp Bounds for Some Multicolor Ramsey Numbers, *Combinatorica*, **25** (2005) 125-141.
- [-] B.M.N. Alzaleq, see [JarAl].
- [AKM] J. Arste, K. Klamroth and I. Mengersen, Three Color Ramsey Numbers for Small Graphs, *Utilitas Mathematica*, **49** (1996) 85-96.
- [-] H. Assiyatun, see [HaBA1, HaBA2, BaHA].
- [AFM] M. Axenovich, Z. Füredi and D. Mubayi, On Generalized Ramsey Theory: the Bipartite Case, *Journal of Combinatorial Theory, Series B*, **79** (2000) 66-86.

B

- [BaRT]* A. Babak, S.P. Radziszowski and Kung-Kuen Tse, Computation of the Ramsey Number $R(B_3, K_5)$, *Bulletin of the Institute of Combinatorics and its Applications*, **41** (2004) 71-76.
- [BaiLi] Bai Lufeng and Li Yusheng, Algebraic Constructions and Applications in Ramsey Theory, *Advances in Mathematics*, **35** (2006) 167-170.
- [BaLX] Bai Lufeng, Li Yusheng and Xu Zhiqiang, Algebraic Constructions and Applications in Ramsey Theory, *Journal of Mathematical Study*, **37** (2004) 245-249.

- [–] Bai Lufeng, see also [SoBL].
- [BaLS] P.N. Balister, J. Lehel and R.H. Schelp, Ramsey Unsaturated and Saturated Graphs, *Journal of Graph Theory*, **51** (2006) 22-32.
- [BaSS] P.N. Balister, R.H. Schelp and M. Simonovits, A Note on Ramsey Size-Linear Graphs, *Journal of Graph Theory*, **39** (2002) 1-5.
- [–] A.M.M. Baniabedaruhan, see [JarBa].
- [–] Qiquan Bao, see [ShaXBP].
- [Bas] E.T. Baskoro, The Ramsey Number of Paths and Small Wheels, *Majalah Ilmiah Himpunan Matematika Indonesia*, MIHMI, **8** (2002) 13-16.
- [BaHA] E.T. Baskoro, Hasmawati and H. Assiyatun, The Ramsey Numbers for Disjoint Unions of Trees, *Discrete Mathematics*, **306** (2006) 3297-3301.
- [BaSu] E.T. Baskoro and Surahmat, The Ramsey Number of Paths with respect to Wheels, *Discrete Mathematics*, **294** (2005) 275-277.
- [BSNM] E.T. Baskoro, Surahmat, S.M. Nababan and M. Miller, On Ramsey Graph Numbers for Trees versus Wheels of Five or Six Vertices, *Graphs and Combinatorics*, **18** (2002) 717-721.
- [–] E.T. Baskoro, see also [AliBas, AliBT, HaBA1, HaBA2, SuBa1, SuBa2, SuBB1, SuBB2, SuBB3, SuBB4, SuBT1, SuBT2, SuBTB, SuBUB].
- [Bev] D. Bevan, *personal communication* (2002).
- [BePi] A. Beveridge and O. Pikhurko, On the Connectivity of Extremal Ramsey Graphs, *Australasian Journal of Combinatorics*, **41** (2008) 57-61.
- [BS] A. Bialostocki and J. Schönheim, On Some Turán and Ramsey Numbers for C_4 , in *Graph Theory and Combinatorics* (ed. B. Bollobás), Academic Press, London, (1984) 29-33.
- [Biel1] H. Bielak, Ramsey and 2-local Ramsey Numbers for Disjoint Unions of Cycles, *Discrete Mathematics*, **307** (2007) 319-330.
- [Biel2] H. Bielak, Ramsey Numbers for a Disjoint Union of Some Graphs, *Applied Mathematics Letters*, **22** (2009) 475-477.
- [Bier] J. Bierbrauer, Ramsey Numbers for the Path with Three Edges, *European Journal of Combinatorics*, **7** (1986) 205-206.
- [BB] J. Bierbrauer and A. Brandis, On Generalized Ramsey Numbers for Trees, *Combinatorica*, **5** (1985) 95-107.
- [BLR]* K. Black, D. Leven and S.P. Radziszowski, New Bounds on Some Ramsey Numbers, *in preparation*, (2009).
- [BJYHRZ] B. Bollobás, C.J. Jayawardene, Yang Jian Sheng, Huang Yi Ru, C.C. Rousseau, and Zhang Ke Min, On a Conjecture Involving Cycle-Complete Graph Ramsey Numbers, *Australasian Journal of Combinatorics*, **22** (2000) 63-71.
- [BH] R. Bolze and H. Harborth, The Ramsey Number $r(K_4-x, K_5)$, in *The Theory and Applications of Graphs*, (Kalamazoo, MI, 1980), John Wiley & Sons, New York, (1981) 109-116.
- [BoEr] J.A. Bondy and P. Erdős, Ramsey Numbers for Cycles in Graphs, *Journal of Combinatorial Theory, Series B*, **14** (1973) 46-54.
- [–] A. Brandis, see [BB].
- [Bra1] S. Brandt, Subtrees and Subforests in Graphs, *Journal of Combinatorial Theory, Series B*, **61** (1994) 63-70.
- [Bra2] S. Brandt, Sufficient Conditions for Graphs to Contain All Subgraphs of a Given Type, *Ph.D. thesis*, Freie Universität Berlin, 1994.
- [Bra3] S. Brandt, Expanding Graphs and Ramsey Numbers, *manuscript*, (1996).
- [BBH1]** S. Brandt, G. Brinkmann and T. Harmuth, All Ramsey Numbers $r(K_3, G)$ for Connected Graphs of Order 9, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R7, **5** (1998), 20

pages.

- [BBH2]** S. Brandt, G. Brinkmann and T. Harmuth, The Generation of Maximal Triangle-Free Graphs, *Graphs and Combinatorics*, **16** (2000) 149-157.
- [Brin]** G. Brinkmann, All Ramsey Numbers $r(K_3, G)$ for Connected Graphs of Order 7 and 8, *Combinatorics, Probability and Computing*, **7** (1998) 129-140.
- [-] G. Brinkmann, see also [BBH1, BBH2].
- [-] H.J. Broersma, see [SaBr1, SaBr2, SaBr3, SaBr4, SuBB1, SuBB2, SuBB3, SuBB4, SuBTB, SuBUB].
- [BR]* J.P. Burling and S.W. Reyner, Some Lower Bounds of the Ramsey Numbers $n(k, k)$, *Journal of Combinatorial Theory, Series B*, **13** (1972) 168-169.
- [Bu1] S.A. Burr, Generalized Ramsey Theory for Graphs - a Survey, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 52-75.
- [Bu2] S.A. Burr, Ramsey Numbers Involving Graphs with Long Suspended Paths, *Journal of the London Mathematical Society* (2), **24** (1981) 405-413.
- [Bu3] S.A. Burr, Multicolor Ramsey Numbers Involving Graphs with Long Suspended Path, *Discrete Mathematics*, **40** (1982) 11-20.
- [Bu4] S.A. Burr, Diagonal Ramsey Numbers for Small Graphs, *Journal of Graph Theory*, **7** (1983) 57-69.
- [Bu5] S.A. Burr, Ramsey Numbers Involving Powers of Sparse Graphs, *Ars Combinatoria*, **15** (1983) 163-168.
- [Bu6] S.A. Burr, Determining Generalized Ramsey Numbers is NP-Hard, *Ars Combinatoria*, **17** (1984) 21-25.
- [Bu7] S.A. Burr, What Can We Hope to Accomplish in Generalized Ramsey Theory?, *Discrete Mathematics*, **67** (1987) 215-225.
- [Bu8] S.A. Burr, On the Ramsey Numbers $r(G, nH)$ and $r(nG, nH)$ When n Is Large, *Discrete Mathematics*, **65** (1987) 215-229.
- [Bu9] S.A. Burr, On Ramsey Numbers for Large Disjoint Unions of Graphs, *Discrete Mathematics*, **70** (1988) 277-293.
- [Bu10] S.A. Burr, On the Computational Complexity of Ramsey-type Problems, Mathematics of Ramsey Theory, *Algorithms and Combinatorics*, **5**, Springer, Berlin, 1990, 46-52.
- [BE1] S.A. Burr and P. Erdős, On the Magnitude of Generalized Ramsey Numbers for Graphs, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds., Keszthely 1973) Vol. 1, 215-240, Colloq. Math. Soc. Janos Bolyai, Vol. 10, North-Holland, Amsterdam, 1975.
- [BE2] S.A. Burr and P. Erdős, Extremal Ramsey Theory for Graphs, *Utilitas Mathematica*, **9** (1976) 247-258.
- [BE3] S.A. Burr and P. Erdős, Generalizations of a Ramsey-Theoretic Result of Chvátal, *Journal of Graph Theory*, **7** (1983) 39-51.
- [BEFRS1] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, An Extremal Problem in Generalized Ramsey Theory, *Ars Combinatoria*, **10** (1980) 193-203.
- [BEFRS2] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Numbers for the Pair Sparse Graph-Path or Cycle, *Transactions of the American Mathematical Society*, **269** (1982) 501-512.
- [BEFRS3] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, On Ramsey Numbers Involving Starlike Multipartite Graphs, *Journal of Graph Theory*, **7** (1983) 395-409.
- [BEFRS4] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, The Ramsey Number for the Pair Complete Bipartite Graph-Graph of Limited Degree, in *Graph Theory with Applications to Algorithms and Computer Science*, (Y. Alavi et al. eds.), John Wiley & Sons, New York, (1985) 163-174.
- [BEFRS5] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Some Complete Bipartite Graph-Tree Ramsey Numbers, *Annals of Discrete Mathematics*, **41** (1989) 79-89.

- [BEFRSGJ] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau, R.H. Schelp, R.J. Gould and M.S. Jacobson, Goodness of Trees for Generalized Books, *Graphs and Combinatorics*, **3** (1987) 1-6.
- [BEFS] S.A. Burr, P. Erdős, R.J. Faudree and R.H. Schelp, On the Difference between Consecutive Ramsey Numbers, *Utilitas Mathematica*, **35** (1989) 115-118.
- [BES] S.A. Burr, P. Erdős and J.H. Spencer, Ramsey Theorems for Multiple Copies of Graphs, *Transactions of the American Mathematical Society*, **209** (1975) 87-99.
- [BF] S.A. Burr and R.J. Faudree, On Graphs G for Which All Large Trees Are G -good, *Graphs and Combinatorics*, **9** (1993) 305-313.
- [BG] S.A. Burr and J.W. Grossman, Ramsey Numbers of Graphs with Long Tails, *Discrete Mathematics*, **41** (1982) 223-227.
- [BuRo1] S.A. Burr and J.A. Roberts, On Ramsey Numbers for Stars, *Utilitas Mathematica*, **4** (1973) 217-220.
- [BuRo2] S.A. Burr and J.A. Roberts, On Ramsey Numbers for Linear Forests, *Discrete Mathematics*, **8** (1974) 245-250.
- [BuRo3] S.A. Burr and V. Rosta, On the Ramsey Multiplicities of Graphs - Problems and Recent Results, *Journal of Graph Theory*, **4** (1980) 347-361.
- [Bush] L.E. Bush, The William Lowell Putnam Mathematical Competition (question #2 in Part I asks for the proof of $R(3,3) \leq 6$), *American Mathematical Monthly*, **60** (1953) 539-542.

C

- [CET]* N.J. Calkin, P. Erdős and C.A. Tovey, New Ramsey Bounds from Cyclic Graphs of Prime Order, *SIAM Journal of Discrete Mathematics*, **10** (1997) 381-387.
- [Car] D. Cariolaro, On the Ramsey Number $R(3, 6)$, *Australasian J. of Combinatorics*, **37** (2007) 301-304.
- [Caro] Y. Caro, Zero-Sum Problems - A Survey, *Discrete Mathematics*, **152** (1996) 93-113.
- [CLRZ] Y. Caro, Li Yusheng, C.C. Rousseau and Zhang Yuming, Asymptotic Bounds for Some Bipartite Graph - Complete Graph Ramsey Numbers, *Discrete Mathematics*, **220** (2000) 51-56.
- [CGP] G. Chartrand, R.J. Gould and A.D. Polimeni, On Ramsey Numbers of Forests versus Nearly Complete Graphs, *Journal of Graph Theory*, **4** (1980) 233-239.
- [CRSPS] G. Chartrand, C.C. Rousseau, M.J. Stewart, A.D. Polimeni and J. Sheehan, On Star-Book Ramsey Numbers, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 203-214.
- [ChaS] G. Chartrand and S. Schuster, On the existence of specified cycles in complementary graphs, *Bulletin of the American Mathematical Society*, **77** (1971) 995-998.
- [Chen] Chen Guantao, A Result on C_4 -Star Ramsey Numbers, *Discrete Mathematics*, **163** (1997) 243-246.
- [ChenS] Chen Guantao and R.H. Schelp, Graphs with Linearly Bounded Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **57** (1993) 138-149.
- [-] Chen Hong, see [WuCXS].
- [ChenJ] Chen Jie, The Lower Bound of Some Ramsey Numbers (in Chinese), *Journal of Liaoning Normal University, Natural Science*, **25** (2002) 244-246.
- [ChenCMN] Yaojun Chen, T.C. Edwin Cheng, Zhengke Miao and C.T. Ng, The Ramsey Numbers for Cycles versus Wheels of Odd Order, *to appear*, (2009).
- [ChenCN] Yaojun Chen, T.C. Edwin Cheng and C.T. Ng, A Theorem on Cycle-Wheel Ramsey Number, *manuscript*, (2009).
- [ChenCX] Yaojun Chen, T.C. Edwin Cheng and Ran Xu, The Ramsey Number for a Cycle of Length Six versus a Clique of Order Eight, *Discrete Applied Mathematics*, **157** (2009) 8-12.
- [ChenCZ1] Yaojun Chen, T.C. Edwin Cheng and Yunqing Zhang, The Ramsey Numbers $R(C_m, K_7)$ and $R(C_7, K_8)$, *European Journal of Combinatorics*, **29** (2008) 1337-1352.

- [ChenCZ2] Yaojun Chen, T.C. Edwin Cheng and Yunqing Zhang, The Ramsey Numbers for Stars of Odd Order versus a Wheel of Order Nine, *to appear*, (2009).
- [ChenZZ1] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Paths versus Wheels, *Discrete Mathematics*, **290** (2005) 85-87.
- [ChenZZ2] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Stars versus Wheels, *European Journal of Combinatorics*, **25** (2004) 1067-1075.
- [ChenZZ3] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers $R(T_n, W_6)$ for $\Delta(T_n) \geq n - 3$, *Applied Mathematics Letters*, **17** (2004) 281-285.
- [ChenZZ4] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Trees versus W_6 or W_7 , *European Journal of Combinatorics*, **27** (2006) 558-564.
- [ChenZZ5] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers $R(T_n, W_6)$ for Small n , *Utilitas Mathematica*, **67** (2005) 269-284.
- [ChenZZ6] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers $R(T_n, W_6)$ for T_n without Certain Deletable Sets, *Journal of Systems Science and Complexity*, **18** (2005) 95-101.
- [-] Chen Yaojun, see also [CheCZN, ZhaCC, ZhaCZ1, ZhaCZ2].
- [Cheng] Cheng Ying, On Graphs Which Do Not Contain Certain Trees, *Ars Combinatoria*, **19** (1985) 119-151.
- [CheCZN] T.C. Edwin Cheng, Yaojun Chen, Yunqing Zhang and C.T. Ng, The Ramsey Numbers for a Cycle of Length Six or Seven versus a Clique of Order Seven, *Discrete Mathematics*, **307** (2007) 1047-1053.
- [-] T.C. Edwin Cheng, see also [ChenCMN, ChenCN, ChenCX, ChenCZ1, ChenCZ2, ZhaCC].
- [Chu1] F.R.K. Chung, On the Ramsey Numbers $N(3,3,\dots,3; 2)$, *Discrete Mathematics*, **5** (1973) 317-321.
- [Chu2] F.R.K. Chung, On Triangular and Cyclic Ramsey Numbers with k Colors, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 236-241.
- [Chu3] F.R.K. Chung, A Note on Constructive Methods for Ramsey Numbers, *Journal of Graph Theory*, **5** (1981) 109-113.
- [Chu4] F.R.K. Chung, Open problems of Paul Erdős in Graph Theory, *Journal of Graph Theory*, **25** (1997) 3-36.
- [CCD] F.R.K. Chung, R. Cleve and P. Dagum, A Note on Constructive Lower Bounds for the Ramsey Numbers $R(3, t)$, *Journal of Combinatorial Theory, Series B*, **57** (1993) 150-155.
- [ChGra1] F.R.K. Chung and R.L. Graham, On Multicolor Ramsey Numbers for Complete Bipartite Graphs, *Journal of Combinatorial Theory, Series B*, **18** (1975) 164-169.
- [ChGra2] F.R.K. Chung and R.L. Graham, *Erdős on Graphs, His Legacy of Unsolved Problems*, A K Peters, Wellesley, Massachusetts (1998).
- [ChGri] F.R.K. Chung and C.M. Grinstead, A Survey of Bounds for Classical Ramsey Numbers, *Journal of Graph Theory*, **7** (1983) 25-37.
- [Chv] V. Chvátal, Tree-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **1** (1977) 93.
- [CH1] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, II. Small Diagonal Numbers, *Proceedings of the American Mathematical Society*, **32** (1972) 389-394.
- [CH2] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, III. Small Off-Diagonal Numbers, *Pacific Journal of Mathematics*, **41** (1972) 335-345.
- [CH3] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, I. Diagonal Numbers, *Periodica Mathematica Hungarica*, **3** (1973) 115-124.
- [CRST] V. Chvátal, V. Rödl, E. Szemerédi and W.T. Trotter Jr., The Ramsey Number of a Graph with Bounded Maximum Degree, *Journal of Combinatorial Theory, Series B*, **34** (1983) 239-243.
- [ChvS] V. Chvátal and A. Schwenk, On the Ramsey Number of the Five-Spoked Wheel, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 247-261.

- [Clan] M. Clancy, Some Small Ramsey Numbers, *Journal of Graph Theory*, **1** (1977) 89-91.
- [Clap] C. Clapham, The Ramsey Number $r(C_4, C_4, C_4)$, *Periodica Mathematica Hungarica*, **18** (1987) 317-318.
- [CEHMS] C. Clapham, G. Exoo, H. Harborth, I. Mengersen and J. Sheehan, The Ramsey Number of $K_5 - e$, *Journal of Graph Theory*, **13** (1989) 7-15.
- [Clark] L. Clark, On Cycle-Star Graph Ramsey Numbers, *Congressus Numerantium*, **50** (1985) 187-192.
- [-] R. Cleve, see [CCD].
- [Coc] E.J. Cockayne, Some Tree-Star Ramsey Numbers, *Journal of Combinatorial Theory*, Series B, **17** (1974) 183-187.
- [CL1] E.J. Cockayne and P.J. Lorimer, The Ramsey Number for Stripes, *Journal of the Australian Mathematical Society*, Series A, **19** (1975) 252-256.
- [CL2] E.J. Cockayne and P.J. Lorimer, On Ramsey Graph Numbers for Stars and Stripes, *Canadian Mathematical Bulletin*, **18** (1975) 31-34.
- [CPR] B. Codenotti, P. Pudlák and G. Resta, Some Structural Properties of Low-Rank Matrices Related to Computational Complexity, *Theoretical Computer Science*, **235** (2000) 89-107.
- [CoPC] Special issue on Ramsey theory of *Combinatorics, Probability and Computing*, **12** (2003), Numbers 5 and 6.
- [Con1] D. Conlon, A New Upper Bound for Diagonal Ramsey Numbers, to appear in the *Annals of Mathematics*.
- [Con2] D. Conlon, Hypergraph Packing and Sparse Bipartite Ramsey Numbers, to appear in *Combinatorics, Probability and Computing*.
- [ConFS1] D. Conlon, J. Fox and B. Sudakov, Hypergraph Ramsey Numbers, *manuscript*, (2009).
- [ConFS2] D. Conlon, J. Fox and B. Sudakov, Large Almost Monochromatic Subsets in Hypergraphs, *manuscript*, (2009).
- [ConFS3] D. Conlon, J. Fox and B. Sudakov, Ramsey Numbers of Sparse Hypergraphs, *Random Structures and Algorithms*, **35** (2009) 1-14.
- [CooFKO] O. Cooley, N. Fountoulakis, D. Kühn and D. Osthus, 3-Uniform Hypergraphs of Bounded Degree Have Linear Ramsey Numbers, *Journal of Combinatorial Theory*, Series B, **98** (2008) 484-505.
- [-] O. Cooley, see also [KüCFO].
- [CsKo] R. Csákány and J. Komlós, The Smallest Ramsey Numbers, *Discrete Mathematics*, **199** (1999) 193-199.

D

- [-] P. Dagum, see [CCD].
- [Den] T. Denley, The Ramsey Numbers for Disjoint Unions of Cycles, *Discrete Mathematics*, **149** (1996) 31-44.
- [DoLL] Lin Dong, Yusheng Li and Qizhong Lin, Ramsey Numbers Involving Graphs with Large Degrees, to appear in *Applied Mathematics Letters*, (2009).
- [DuHu] Duan Chanlun and Huang Wenke, Lower Bound of Ramsey Number $r(3, 10)$ (in Chinese), *Acta Scientiarum Naturalium Universitatis Nei Mongol*, **31** (2000) 468-470.
- [DLR] D. Duffus, H. Lefmann and V. Rödl, Shift Graphs and Lower Bounds on Ramsey Numbers $r_k(l; r)$, *Discrete Mathematics*, **137** (1995) 177-187.
- [Dzi]* T. Dzido, Multicolor Ramsey Numbers for Paths and Cycles, *Discussiones Mathematicae Graph Theory*, **25** (2005) 57-65.
- [DzFi1]* T. Dzido and R. Fidytek, The Number of Critical Colorings for Some Ramsey Numbers, *International Journal of Pure and Applied Mathematics*, ISSN 1311-8080, **38** (2007) 433-444.

- [DzFi2]* T. Dzido and R. Fidytek, On Some Three Color Ramsey Numbers for Paths and Cycles, *Discrete Mathematics*, **309** (2009) 4955-4958.
- [DzKP] T. Dzido, M. Kubale and K. Piwakowski, On Some Ramsey and Turán-type Numbers for Paths and Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R55, **13** (2006), 9 pages.
- [DzNS] T. Dzido, A. Nowik and P. Szuca, New Lower Bound for Multicolor Ramsey Numbers for Even Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #N13, **12** (2005), 5 pages.

E

- [Ea1] Easy to obtain by simple combinatorics from other results, in particular by using graphs establishing lower bounds with smaller parameters.
- [Ea2] Unique 2-(6,3,2) design gives lower bound 7, upper bound is easy.
- [Eaton] N. Eaton, Ramsey Numbers for Sparse Graphs, *Discrete Mathematics*, **185** (1998) 63-75.
- [Erd1] P. Erdős, Some Remarks on the Theory of Graphs, *Bulletin of the American Mathematical Society*, **53** (1947) 292-294.
- [Erd2] P. Erdős, On the Combinatorial Problems Which I Would Most Like to See Solved, *Combinatorica*, **1** (1981) 25-42.
- [EFRS1] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Generalized Ramsey Theory for Multiple Colors, *Journal of Combinatorial Theory, Series B*, **20** (1976) 250-264.
- [EFRS2] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, On Cycle-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **2** (1978) 53-64.
- [EFRS3] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Numbers for Brooms, *Congressus Numerantium*, **35** (1982) 283-293.
- [EFRS4] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Multipartite Graph-Sparse Graph Ramsey Numbers, *Combinatorica*, **5** (1985) 311-318.
- [EFRS5] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, A Ramsey Problem of Harary on Graphs with Prescribed Size, *Discrete Mathematics*, **67** (1987) 227-233.
- [EFRS6] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Extremal Theory and Bipartite Graph-Tree Ramsey Numbers, *Discrete Mathematics*, **72** (1988) 103-112.
- [EFRS7] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, The Book-Tree Ramsey Numbers, *Scientia, Series A: Mathematical Sciences, Valparaíso, Chile*, **1** (1988) 111-117.
- [EFRS8] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Multipartite Graph-Tree Graph Ramsey Numbers, in *Graph Theory and Its Applications: East and West, Proceedings of the First China-USA International Graph Theory Conference*, Annals of the New York Academy of Sciences, **576** (1989) 146-154.
- [EFRS9] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Size Linear Graphs, *Combinatorics, Probability and Computing*, **2** (1993) 389-399.
- [EG] P. Erdős and R.L. Graham, On Partition Theorems for Finite Sets, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds.) Vol. 1, 515--527, Colloq. Math. Soc. Janos Bolyai, Vol. 10, North Holland, 1975.
- [-] P. Erdős, see also [BoEr, BE1, BE2, BE3, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, BEFS, BES, CET].
- [Ex1]* G. Exoo, Ramsey Numbers of Hypergraphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **2** (1987) 5-11.
- [Ex2]* G. Exoo, Constructing Ramsey Graphs with a Computer, *Congressus Numerantium*, **59** (1987) 31-36.

- [Ex3]* G. Exoo, Applying Optimization Algorithm to Ramsey Problems, in *Graph Theory, Combinatorics, Algorithms, and Applications* (Y. Alavi ed.), SIAM Philadelphia, (1989) 175-179.
- [Ex4]* G. Exoo, A Lower Bound for $R(5, 5)$, *Journal of Graph Theory*, **13** (1989) 97-98.
- [Ex5]* G. Exoo, On Two Classical Ramsey Numbers of the Form $R(3, n)$, *SIAM Journal of Discrete Mathematics*, **2** (1989) 488-490.
- [Ex6]* G. Exoo, A Lower Bound for $r(K_5 - e, K_5)$, *Utilitas Mathematica*, **38** (1990) 187-188.
- [Ex7]* G. Exoo, On the Three Color Ramsey Number of $K_4 - e$, *Discrete Mathematics*, **89** (1991) 301-305.
- [Ex8]* G. Exoo, Indiana State University, *personal communication* (1992).
- [Ex9]* G. Exoo, Announcement: On the Ramsey Numbers $R(4, 6)$, $R(5, 6)$ and $R(3, 12)$, *Ars Combinatoria*, **35** (1993) 85. The construction of a graph proving $R(4, 6) \geq 35$ is presented in detail at <http://ginger.indstate.edu/ge/RAMSEY> (2001).
- [Ex10]* G. Exoo, A Lower Bound for Schur Numbers and Multicolor Ramsey Numbers of K_3 , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R8, **1** (1994), 3 pages.
- [Ex11]* G. Exoo, Indiana State University, *personal communication* (1997).
- [Ex12]* G. Exoo, Some New Ramsey Colorings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R29, **5** (1998), 5 pages. The constructions are available electronically from <http://ginger.indstate.edu/ge/RAMSEY>.
- [Ex13]* G. Exoo, Indiana State University, *personal communication* (1998). Constructions available at <http://ginger.indstate.edu/ge/RAMSEY>.
- [Ex14]* G. Exoo, Indiana State University, *New Lower Bounds for Table III*, (2000). Constructions available at <http://ginger.indstate.edu/ge/RAMSEY>.
- [Ex15]* G. Exoo, Some Applications of pq -groups in Graph Theory, *Discussiones Mathematicae Graph Theory*, **24** (2004) 109-114. Constructions available at <http://ginger.indstate.edu/ge/RAMSEY>.
- [Ex16]* G. Exoo, Indiana State University, *personal communication* (2002-2004). Constructions available at <http://ginger.indstate.edu/ge/RAMSEY>.
- [Ex17]* G. Exoo, Indiana State University, *personal communication* (2005-2006). Constructions available at <http://ginger.indstate.edu/ge/RAMSEY>.
- [EHM1] G. Exoo, H. Harborth and I. Mengersen, The Ramsey Number of K_4 versus $K_5 - e$, *Ars Combinatoria*, **25A** (1988) 277-286.
- [EHM2] G. Exoo, H. Harborth and I. Mengersen, On Ramsey Number of $K_{2,n}$, in *Graph Theory, Combinatorics, Algorithms, and Applications* (Y. Alavi, F.R.K. Chung, R.L. Graham and D.F. Hsu eds.), SIAM Philadelphia, (1989) 207-211.
- [ExRe]* G. Exoo and D.F. Reynolds, Ramsey Numbers Based on C_5 -Decompositions, *Discrete Mathematics*, **71** (1988) 119-127.
- [-] G. Exoo, see also [CEHMS, XXER].

F

- [FLPS] R.J. Faudree, S.L. Lawrence, T.D. Parsons and R.H. Schelp, Path-Cycle Ramsey Numbers, *Discrete Mathematics*, **10** (1974) 269-277.
- [FM]** R.J. Faudree and B.D. McKay, A Conjecture of Erdős and the Ramsey Number $r(W_6)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **13** (1993) 23-31.
- [FRS1] R.J. Faudree, C.C. Rousseau and R.H. Schelp, All Triangle-Graph Ramsey Numbers for Connected Graphs of Order Six, *Journal of Graph Theory*, **4** (1980) 293-300.
- [FRS2] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Studies Related to the Ramsey Number $r(K_5 - e)$, in *Graph Theory and Its Applications to Algorithms and Computer Science*, (Y. Alavi et al. eds.), John Wiley and Sons, New York, (1985) 251-271.

- [FRS3] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Generalizations of the Tree-Complete Graph Ramsey Number, in *Graphs and Applications*, (F. Harary and J.S. Maybee eds.), John Wiley and Sons, New York, (1985) 117-126.
- [FRS4] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Small Order Graph-Tree Ramsey Numbers, *Discrete Mathematics*, **72** (1988) 119-127.
- [FRS5] R.J. Faudree, C.C. Rousseau and R.H. Schelp, A Good Idea in Ramsey Theory, in *Graph Theory, Combinatorics, Algorithms, and Applications* (San Francisco, CA 1989), SIAM Philadelphia, PA (1991) 180-189.
- [FRS6] R.J. Faudree, C.C. Rousseau and J. Sheehan, More from the Good Book, in *Proceedings of the Ninth Southeastern Conference on Combinatorics, Graph Theory, and Computing*, Utilitas Mathematica Publ., *Congressus Numerantium XXI* (1978) 289-299.
- [FRS7] R.J. Faudree, C.C. Rousseau and J. Sheehan, Strongly Regular Graphs and Finite Ramsey Theory, *Linear Algebra and its Applications*, **46** (1982) 221-241.
- [FRS8] R.J. Faudree, C.C. Rousseau and J. Sheehan, Cycle-Book Ramsey Numbers, *Ars Combinatoria*, **31** (1991) 239-248.
- [FS1] R.J. Faudree and R.H. Schelp, All Ramsey Numbers for Cycles in Graphs, *Discrete Mathematics*, **8** (1974) 313-329.
- [FS2] R.J. Faudree and R.H. Schelp, Path Ramsey Numbers in Multicolorings, *Journal of Combinatorial Theory, Series B*, **19** (1975) 150-160.
- [FS3] R.J. Faudree and R.H. Schelp, Ramsey Numbers for All Linear Forests, *Discrete Mathematics*, **16** (1976) 149-155.
- [FS4] R.J. Faudree and R.H. Schelp, Some Problems in Ramsey Theory, in *Theory and Applications of Graphs*, (conference proceedings, Kalamazoo, MI 1976), Lecture Notes in Mathematics **642**, Springer, Berlin, (1978) 500-515.
- [FSR] R.J. Faudree, R.H. Schelp and C.C. Rousseau, Generalizations of a Ramsey Result of Chvátal, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 351-361.
- [FSS1] R.J. Faudree, R.H. Schelp and M. Simonovits, On Some Ramsey Type Problems Connected with Paths, Cycles and Trees, *Ars Combinatoria*, **29A** (1990) 97-106.
- [FSS2] R.J. Faudree, A. Schelten and I. Schiermeyer, The Ramsey Number $r(C_7, C_7, C_7)$, *Discussiones Mathematicae Graph Theory*, **23** (2003) 141-158.
- [FS] R.J. Faudree and M. Simonovits, Ramsey Problems and Their Connection to Turán-Type Extremal Problems, *Journal of Graph Theory*, **16** (1992) 25-50.
- [-] R.J. Faudree, see also [BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, BEFS, BF, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9].
- [FKR]** S. Fettes, R.L. Kramer and S.P. Radziszowski, An Upper Bound of 62 on the Classical Ramsey Number $R(3,3,3,3)$, *Ars Combinatoria*, **72** (2004) 41-63.
- [-] R. Fidytek, see [DzFi1, DzFi2].
- [FiŁu] A. Figaj and T. Łuczak, The Ramsey Number for a Triple of Long Even Cycles, *Journal of Combinatorial Theory, Series B*, **97** (2007) 584-596.
- [Fo] J. Folkman, Notes on the Ramsey Number $N(3,3,3,3)$, *Journal of Combinatorial Theory, Series A*, **16** (1974) 371-379.
- [-] N. Fountoulakis, see [CooFKO, KüCFO].
- [FoxSu1] J. Fox and B. Sudakov, Density Theorems for Bipartite Graphs and Related Ramsey-type Results, *to appear*, (2009).
- [FoxSu2] J. Fox and B. Sudakov, Two Remarks on the Burr-Erdős Conjecture, *to appear*, (2009).
- [-] J. Fox, see also [ConFS1, ConFS2, ConFS3].

- [FraWi] P. Frankl and R.M. Wilson, Intersection Theorems with Geometric Consequences, *Combinatorica*, **1** (1981) 357-368.
- [Fra1] K. Fraughnaugh Jones, Independence in Graphs with Maximum Degree Four, *Journal of Combinatorial Theory, Series B*, **37** (1984) 254-269.
- [Fra2] K. Fraughnaugh Jones, Size and Independence in Triangle-Free Graphs with Maximum Degree Three, *Journal of Graph Theory*, **14** (1990) 525-535.
- [FrLo] K. Fraughnaugh and S.C. Locke, Finding Independent Sets in Triangle-free Graphs, *SIAM Journal of Discrete Mathematics*, **9** (1996) 674-681.
- [Fre] H. Fredricksen, Schur Numbers and the Ramsey Numbers $N(3,3,\dots,3;2)$, *Journal of Combinatorial Theory, Series A*, **27** (1979) 376-377.
- [FreSw]* H. Fredricksen and M.M. Sweet, Symmetric Sum-Free Partitions and Lower Bounds for Schur Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R32, **7** (2000), 9 pages.
- [-] Z. Füredi, see [AFM].

G

- [Gas] W. Gasarch, Applications of Ramsey Theory to Computer Science, collection of pointers to papers, <http://www.cs.umd.edu/~gasarch/ramsey/ramsey.html> (2009).
- [GauST] S. Gautam, A.K. Srivastava and A. Tripathi, On Multicolour Noncomplete Ramsey Graphs of Star Graphs, *Discrete Applied Mathematics*, **156** (2008) 2423-2428.
- [GeGy] L. Gerencsér and A. Gyárfás, On Ramsey-Type Problems, *Annales Universitatis Scientiarum Budapestinensis, Eötvös Sect. Math.*, **10** (1967) 167-170.
- [Gi1] G. Giraud, Une généralisation des nombres et de l'inégalité de Schur, *C.R. Acad. Sc. Paris, Séries A-B*, **266** (1968) A437-A440.
- [Gi2] G. Giraud, Minoration de certains nombres de Ramsey binaires par les nombres de Schur généralisés, *C.R. Acad. Sc. Paris, Séries A-B*, **266** (1968) A481-A483.
- [Gi3] G. Giraud, Nouvelles majorations des nombres de Ramsey binaires-bicolores, *C.R. Acad. Sc. Paris, Séries A-B*, **268** (1969) A5-A7.
- [Gi4] G. Giraud, Majoration du nombre de Ramsey ternaire-bicolore en (4,4), *C.R. Acad. Sc. Paris, Séries A-B*, **269** (1969) A620-A622.
- [Gi5] G. Giraud, Une minoration du nombre de quadrangles unicolores et son application à la majoration des nombres de Ramsey binaires-bicolores, *C.R. Acad. Sc. Paris, Séries A-B*, **276** (1973) A1173-A1175.
- [Gi6] G. Giraud, Sur le problème de Goodman pour les quadrangles et la majoration des nombres de Ramsey, *Journal of Combinatorial Theory, Series B*, **27** (1979) 237-253.
- [-] A.M. Gleason, see [GG].
- [GK] W. Goddard and D.J. Kleitman, An upper bound for the Ramsey numbers $r(K_3, G)$, *Discrete Mathematics*, **125** (1994) 177-182.
- [GoMC] A. Gonçalves and E.L. Monte Carmelo, Some Geometric Structures and Bounds for Ramsey Numbers, *Discrete Mathematics*, **280** (2004) 29-38.
- [GoJa1] R.J. Gould and M.S. Jacobson, Bounds for the Ramsey Number of a Disconnected Graph Versus Any Graph, *Journal of Graph Theory*, **6** (1982) 413-417.
- [GoJa2] R.J. Gould and M.S. Jacobson, On the Ramsey Number of Trees Versus Graphs with Large Clique Number, *Journal of Graph Theory*, **7** (1983) 71-78.
- [-] R.J. Gould, see also [BEFRSGJ, CGP].
- [GrNe] R.L. Graham and J. Nešetřil, Ramsey Theory and Paul Erdős (Recent Results from a Historical Perspective), *Bolyai Society Mathematical Studies*, **11**, Budapest (2002) 339-365.

- [GrRö] R.L. Graham and V. Rödl, Numbers in Ramsey Theory, in *Surveys in Combinatorics*, (ed. C. Whitehead), Cambridge University Press, 1987, 111-153.
- [GRR1] R.L. Graham, V. Rödl and A. Ruciński, On Graphs with Linear Ramsey Numbers, *Journal of Graph Theory*, **35** (2000) 176-192.
- [GRR2] R.L. Graham, V. Rödl and A. Ruciński, On Bipartite Graphs with Linear Ramsey Numbers, Paul Erdős and his mathematics, *Combinatorica*, **21** (2001) 199-209.
- [GRS] R.L. Graham, B.L. Rothschild and J.H. Spencer, *Ramsey Theory*, John Wiley & Sons, 1990.
- [-] R.L. Graham, see also [ChGra1, ChGra2, EG].
- [GY] J.E. Graver and J. Yackel, Some Graph Theoretic Results Associated with Ramsey's Theorem, *Journal of Combinatorial Theory*, **4** (1968) 125-175.
- [GG] R.E. Greenwood and A.M. Gleason, Combinatorial Relations and Chromatic Graphs, *Canadian Journal of Mathematics*, **7** (1955) 1-7.
- [GH] U. Grenda and H. Harborth, The Ramsey Number $r(K_3, K_7 - e)$, *Journal of Combinatorics, Information & System Sciences*, **7** (1982) 166-169.
- [Gri] J.R. Griggs, An Upper Bound on the Ramsey Numbers $R(3, k)$, *Journal of Combinatorial Theory, Series A*, **35** (1983) 145-153.
- [GR]** C. Grinstead and S. Roberts, On the Ramsey Numbers $R(3, 8)$ and $R(3, 9)$, *Journal of Combinatorial Theory, Series B*, **33** (1982) 27-51.
- [-] C. Grinstead, see also [ChGri].
- [Grol1] V. Grolmusz, Superpolynomial Size Set-Systems with Restricted Intersections mod 6 and Explicit Ramsey Graphs, *Combinatorica*, **20** (2000) 73-88.
- [Grol2] V. Grolmusz, Low Rank Co-Diagonal Matrices and Ramsey Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R15, **7** (2000) 7 pages.
- [Grol3] V. Grolmusz, Set-Systems with Restricted Multiple Intersections, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R8, **9** (2002) 10 pages.
- [Gros1] J.W. Grossman, Some Ramsey Numbers of Unicyclic Graphs, *Ars Combinatoria*, **8** (1979) 59-63.
- [Gros2] J.W. Grossman, The Ramsey Numbers of the Union of Two Stars, *Utilitas Mathematica*, **16** (1979) 271-279.
- [GHK] J.W. Grossman, F. Harary and M. Klawe, Generalized Ramsey Theory for Graphs, X: Double Stars, *Discrete Mathematics*, **28** (1979) 247-254.
- [-] J.W. Grossman, see also [BG].
- [GuLi] Gu Hua and Li Yusheng, On Ramsey Number of $K_{2,t+1}$ vs $K_{1,n}$, *Journal of Nanjing University Mathematical Biquarterly*, **19** (2002) 150-153.
- [GuSL] Gu Hua, Song Hongxue and Liu Xiangyang, Ramsey Numbers $r(K_{1,4}, G)$ for All Three-Partite Graphs G of Order Six, *Journal of Southeast University*, (English Edition), **20** (2004) 378-380.
- [-] Gu Hua, see also [SoGQ].
- [GuoV] Guo Yubao and L. Volkmann, Tree-Ramsey Numbers, *Australasian Journal of Combinatorics*, **11** (1995) 169-175.
- [-] L. Gupta, see [GGS].
- [GGS] S.K. Gupta, L. Gupta and A. Sudan, On Ramsey Numbers for Fan-Fan Graphs, *Journal of Combinatorics, Information & System Sciences*, **22** (1997) 85-93.
- [GyRSS] A. Gyárfás, M. Ruszinkó, G.N. Sárközy and E. Szemerédi, Three-color Ramsey Numbers for Paths, *Combinatorica*, **27** (2007) 35-69. *Corrigendum* in **28** (2008) 499-502.
- [GySS] A. Gyárfás, G.N. Sárközy and E. Szemerédi, The Ramsey Number of Diamond-Matchings and Loose Cycles in Hypergraphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R126, **15(1)** (2008), 14 pages.

- [GyTu] A. Gyárfás and Z. Tuza, An Upper Bound on the Ramsey Number of Trees, *Discrete Mathematics*, **66** (1987) 309-310.
- [-] A. Gyárfás, see also [GeGy].

H

- [Häg] R. Häggkvist, On the Path-Complete Bipartite Ramsey Number, *Discrete Mathematics*, **75** (1989) 243-245.
- [Han]* D. Hanson, Sum-free Sets and Ramsey Numbers, *Discrete Mathematics*, **14** (1976) 57-61.
- [-] D. Hanson, see also [AbbH].
- [Har1] F. Harary, Recent Results on Generalized Ramsey Theory for Graphs, in *Graph Theory and Applications*, (Y. Alavi et al. eds.) Springer, Berlin (1972) 125-138.
- [Har2] F. Harary, Generalized Ramsey Theory I to XIII: Achievement and Avoidance Numbers, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 373-390.
- [-] F. Harary, see also [CH1, CH2, CH3, GHK].
- [HaKr]** H. Harborth and S. Krause, Ramsey Numbers for Circulant Colorings, *Congressus Numerantium*, **161** (2003) 139-150.
- [HaMe1] H. Harborth and I. Mengersen, An Upper Bound for the Ramsey Number $r(K_5-e)$, *Journal of Graph Theory*, **9** (1985) 483-485.
- [HaMe2] H. Harborth and I. Mengersen, All Ramsey Numbers for Five Vertices and Seven or Eight Edges, *Discrete Mathematics*, **73** (1988/89) 91-98.
- [HaMe3] H. Harborth and I. Mengersen, The Ramsey Number of $K_{3,3}$, in *Combinatorics, Graph Theory, and Applications*, Vol. **2** (Y. Alavi, G. Chartrand, O.R. Oellermann and J. Schwenk eds.), John Wiley & Sons, (1991) 639-644.
- [-] H. Harborth, see also [BH, CEHMS, EHM1, EHM2, GH].
- [HaMe4] M. Harborth and I. Mengersen, Some Ramsey Numbers for Complete Bipartite Graphs, *Australasian Journal of Combinatorics*, **13** (1996) 119-128.
- [-] T. Harmuth, see [BBH1, BBH2].
- [HaBA1] Hasmawati, E.T. Baskoro and H. Assiyatun, Star-Wheel Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **55** (2005) 123-128.
- [HaBA2] Hasmawati, E.T. Baskoro and H. Assiyatun, The Ramsey Numbers for Disjoint Unions of Graphs, *Discrete Mathematics*, **308** (2008) 2046-2049.
- [-] Hasmawati, see also [BaHA].
- [HaŁP1+] P.E. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński, M. Simonovits and J. Skokan, The Ramsey Number for Hypergraph Cycles I, *Journal of Combinatorial Theory, Series A*, **113** (2006) 67-83.
- [HaŁP2+] P.E. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński and J. Skokan, The Ramsey Number for 3-Uniform Tight Hypergraph Cycles, *Combinatorics, Probability and Computing*, **18** (2009) 165-203.
- [HaŁT] P.E. Haxell, T. Łuczak and P.W. Tingley, Ramsey Numbers for Trees of Small Maximum Degree, *Combinatorica*, **22** (2002) 287-320.
- [Hein] K. Heinrich, Proper Colourings of K_{15} , *Journal of the Australian Mathematical Society, Series A*, **24** (1977) 465-495.
- [He1] G.R.T. Hendry, Diagonal Ramsey Numbers for Graphs with Seven Edges, *Utilitas Mathematica*, **32** (1987) 11-34.
- [He2] G.R.T. Hendry, Ramsey Numbers for Graphs with Five Vertices, *Journal of Graph Theory*, **13** (1989) 245-248.

- [He3] G.R.T. Hendry, The Ramsey Numbers $r(K_2 + \overline{K}_3, K_4)$ and $r(K_1 + C_4, K_4)$, *Utilitas Mathematica*, **35** (1989) 40-54, addendum in **36** (1989) 25-32.
- [He4] G.R.T. Hendry, Critical Colorings for Clancy's Ramsey Numbers, *Utilitas Mathematica*, **41** (1992) 181-203.
- [He5] G.R.T. Hendry, Small Ramsey Numbers II. Critical Colorings for $r(C_5 + e, K_5)$, *Quaestiones Mathematica*, **17** (1994) 249-258.
- [-] G.R.T. Hendry, see also [YH].
- [HiIr]* R. Hill and R.W. Irving, On Group Partitions Associated with Lower Bounds for Symmetric Ramsey Numbers, *European Journal of Combinatorics*, **3** (1982) 35-50.
- [Hir] J. Hirschfeld, A Lower Bound for Ramsey's Theorem, *Discrete Mathematics*, **32** (1980) 89-91.
- [HoMe] M. Hoeth and I. Mengersen, Ramsey Numbers for Graphs of Order Four versus Connected Graphs of Order Six, *Utilitas Mathematica*, **57** (2000) 3-19.
- [HuSo] Huang Da Ming and Song En Min, Properties and Lower Bounds of the Third Order Ramsey Numbers (in Chinese), *Mathematica Applicata*, **9** (1996) 105-107.
- [Hua1] Huang Guotai, Some Generalized Ramsey Numbers (in Chinese), *Mathematica Applicata*, **1** (1988) 97-101.
- [Hua2] Huang Guotai, An Unsolved Problem of Gould and Jacobson (in Chinese), *Mathematica Applicata*, **9** (1996) 234-236.
- [-] Huang Jian, see [HWSYZH].
- [-] Huang Wenke, see [DuHu].
- [HWSYZH] (also abbreviated by HW+) Huang Yi Ru, Wang Yuandi, Sheng Wancheng, Yang Jiansheng, Zhang Ke Min and Huang Jian, New Upper Bound Formulas with Parameters for Ramsey Numbers, *Discrete Mathematics*, **307** (2007) 760-763.
- [HYZ] Huang Yi Ru, Yang Jian Sheng and Zhang Ke Min, A Note on Ramsey Numbers with Two Parameters, *European Journal of Combinatorics*, **27** (2006) 574-576.
- [HZ1] Huang Yi Ru and Zhang Ke Min, An New Upper Bound Formula for Two Color Classical Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **28** (1998) 347-350.
- [HZ2] Huang Yi Ru and Zhang Ke Min, New Upper Bounds for Ramsey Numbers, *European Journal of Combinatorics*, **19** (1998) 391-394.
- [-] Huang Yi Ru, see also [BJYHRZ, YHZ1, YHZ2].

I

- [Ir] R.W. Irving, Generalised Ramsey Numbers for Small Graphs, *Discrete Mathematics*, **9** (1974) 251-264.
- [-] R.W. Irving, see also [HiIr].
- [Isb1] J.R. Isbell, $N(4,4;3) \geq 13$, *Journal of Combinatorial Theory*, **6** (1969) 210.
- [Isb2] J.R. Isbell, $N(5,4;3) \geq 24$, *Journal of Combinatorial Theory*, Series A, **34** (1983) 379-380.
- [Ishi] Y. Ishigami, Linear Ramsey Numbers for Bounded-Degree Hypergraphs, *Electronic Notes in Discrete Mathematics*, **29** (2007) 47-51.

J

- [Jac] M.S. Jacobson, On the Ramsey Number for Stars and a Complete Graph, *Ars Combinatoria*, **17** (1984) 167-172.
- [-] M.S. Jacobson, see also [BEFRSGJ, GoJa1, GoJa2].

- [JarAl] M.M.M. Jaradat and B.M.N. Alzaleq, The Cycle-Complete Graph Ramsey Number $r(C_8, K_8)$, *SUT Journal of Mathematics*, **43** (2007) 85-98.
- [JarBa] M.M.M. Jaradat and A.M.M. Baniabedlruhman, The Cycle-Complete Graph Ramsey Number $r(C_8, K_7)$, *International Journal of Pure and Applied Mathematics*, **41** (2007) 667-677.
- [JR1] C.J. Jayawardene and C.C. Rousseau, An Upper Bound for the Ramsey Number of a Quadrilateral versus a Complete Graph on Seven Vertices, *Congressus Numerantium*, **130** (1998) 175-188.
- [JR2] C.J. Jayawardene and C.C. Rousseau, Ramsey Numbers $r(C_6, G)$ for All Graphs G of Order Less than Six, *Congressus Numerantium*, **136** (1999) 147-159.
- [JR3] C.J. Jayawardene and C.C. Rousseau, The Ramsey Numbers for a Quadrilateral vs. All Graphs on Six Vertices, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **35** (2000) 71-87. Erratum in **51** (2004) 221.
- [JR4] C.J. Jayawardene and C.C. Rousseau, Ramsey Numbers $r(C_5, G)$ for All Graphs G of Order Six, *Ars Combinatoria*, **57** (2000) 163-173.
- [JR5] C.J. Jayawardene and C.C. Rousseau, The Ramsey Number for a Cycle of Length Five vs. a Complete Graph of Order Six, *Journal of Graph Theory*, **35** (2000) 99-108.
- [-] C.J. Jayawardene, see also [BJYHRZ, RoJa1, RoJa2].
- [-] Jiang Baoqi, see [SunYJLS].
- [Jin]** Jin Xia, Ramsey Numbers Involving a Triangle: Theory & Applications, *Technical Report RIT-TR-93-019*, MS thesis, Department of Computer Science, Rochester Institute of Technology, 1993.
- [-] Jin Xia, see also [RaJi].
- [JGT] Special volume on Ramsey theory of the *Journal of Graph Theory*, **7**, Number 1, (1983).

K

- [Ka1] J.G. Kalbfleisch, Construction of Special Edge-Chromatic Graphs, *Canadian Mathematical Bulletin*, **8** (1965) 575-584.
- [Ka2]* J.G. Kalbfleisch, Chromatic Graphs and Ramsey's Theorem, *Ph.D. thesis*, University of Waterloo, January 1966.
- [Ka3] J.G. Kalbfleisch, On Robillard's Bounds for Ramsey Numbers, *Canadian Mathematical Bulletin*, **14** (1971) 437-440.
- [KaSt] J.G. Kalbfleisch and R.G. Stanton, On the Maximal Triangle-Free Edge-Chromatic Graphs in Three Colors, *Journal of Combinatorial Theory*, **5** (1968) 9-20.
- [KáRos] G. Károlyi and V. Rosta, Generalized and Geometric Ramsey Numbers for Cycles, *Theoretical Computer Science*, **263** (2001) 87-98.
- [KerRo] M. Kerber and C. Rowan, CommonLisp program for computing upper bounds on classical Ramsey numbers, <http://www.cs.bham.ac.uk/~mmk/demos/ramsey-upper-limit.lisp> (2009).
- [Kéry] G. Kéry, On a Theorem of Ramsey (in Hungarian), *Matematikai Lapok*, **15** (1964) 204-224.
- [Kim] J.H. Kim, The Ramsey Number $R(3, t)$ has Order of Magnitude $t^2/\log t$, *Random Structures and Algorithms*, **7** (1995) 173-207.
- [KM1] K. Klamroth and I. Mengersen, Ramsey Numbers of K_3 versus (p, q) -Graphs, *Ars Combinatoria*, **43** (1996) 107-120.
- [KM2] K. Klamroth and I. Mengersen, The Ramsey Number of $r(K_{1,3}, C_4, K_4)$, *Utilitas Mathematica*, **52** (1997) 65-81.
- [-] K. Klamroth, see also [AKM].
- [-] M. Klawe, see [GHK].
- [-] D.J. Kleitman, see [GK].

- [KoSS] Y. Kohayakawa, M. Simonovits and J. Skokan, The 3-colored Ramsey Number of Odd Cycles, *Electronic Notes in Discrete Mathematics*, **19** (2005) 397-402.
- [Köh] W. Köhler, On a Conjecture by Grossman, *Ars Combinatoria*, **23** (1987) 103-106.
- [-] J. Komlós, see [CsKo, AKS].
- [Kor] A. Korolova, Ramsey Numbers of Stars versus Wheels of Similar Sizes, *Discrete Mathematics*, **292** (2005) 107-117.
- [KoRö1] A.V. Kostochka and V. Rödl, On Graphs with Small Ramsey Numbers, *Journal of Graph Theory*, **37** (2001) 198-204.
- [KoRö2] A.V. Kostochka and V. Rödl, On Graphs with Small Ramsey Numbers, II, *Combinatorica*, **24** (2004) 389-401.
- [KoRö3] A.V. Kostochka and V. Rödl, On Ramsey Numbers of Uniform Hypergraphs with Given Maximum Degree, *Journal of Combinatorial Theory, Series A*, **113** (2006) 1555-1564.
- [KoSu] A.V. Kostochka and B. Sudakov, On Ramsey Numbers of Sparse Graphs, *Combinatorics, Probability and Computing*, **12** (2003) 627-641.
- [-] R.L. Kramer, see [FKR].
- [KrRod] I. Krasikov and Y. Roditty, On Some Ramsey Numbers of Unicyclic Graphs, *Bulletin of the Institute of Combinatorics and its Applications*, **33** (2001) 29-34.
- [-] S. Krause, see [HaKr].
- [KLR]* D.L. Kreher, Li Wei and S.P. Radziszowski, Lower Bounds for Multi-Colored Ramsey Numbers From Group Orbits, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 87-95.
- [-] D.L. Kreher, see also [RK1, RK2, RK3, RK4].
- [Kriv] M. Krivelevich, Bounding Ramsey Numbers through Large Deviation Inequalities, *Random Structures and Algorithms*, **7** (1995) 145-155.
- [-] M. Krivelevich, see also [AKS].
- [-] M. Kubale, see [DzKP].
- [KüCFO] D. Kühn, O. Cooley, N. Fountoulakis and D. Osthus, Ramsey Numbers of Sparse Hypergraphs, *Electronic Notes in Discrete Mathematics*, **29** (2007) 29-33.
- [-] D. Kühn, see also [CooFKO].

L

- [-] P.C.B. Lam, see [ShiuLL].
- [La1] S.L. Lawrence, Cycle-Star Ramsey Numbers, *Notices of the American Mathematical Society*, **20** (1973) Abstract A- 420.
- [La2] S.L. Lawrence, Bipartite Ramsey Theory, *Notices of the American Mathematical Society*, **20** (1973) Abstract A- 562.
- [-] S.L. Lawrence, see also [FLPS].
- [LayMa] C. Laywine and J.P. Mayberry, A Simple Construction Giving the Two Non-isomorphic Triangle-Free 3-Colored K_{16} 's, *Journal of Combinatorial Theory, Series B*, **45** (1988) 120-124.
- [LaMu] F. Lazebnik and D. Mubayi, New Lower Bounds for Ramsey Numbers of Graphs and Hypergraphs, *Advances in Applied Mathematics*, **28** (2002) 544-559.
- [LaWo1] F. Lazebnik and A. Woldar, New Lower Bounds on the Multicolor Ramsey Numbers $r_k(C_4)$, *Journal of Combinatorial Theory, Series B*, **79** (2000) 172-176.
- [LaWo2] F. Lazebnik and A. Woldar, General Properties of Some Families of Graphs Defined by Systems of Equations, *Journal of Graph Theory*, **38** (2001) 65-86.

- [Lef] H. Lefmann, Ramsey Numbers for Monotone Paths and Cycles, *Ars Combinatoria*, **35** (1993) 271-279.
- [-] H. Lefmann, see also [DLR].
- [-] J. Lehel, see [BaLS].
- [Les]* A. Lesser, Theoretical and Computational Aspects of Ramsey Theory, *Examensarbeten i Matematik*, Matematiska Institutionen, Stockholms Universitet, **3** (2001).
- [-] D. Leven, see [BLR].
- [-] Li Bingxi, see [SunYWLX, SunYXL].
- [LiWa1] Li Da Yong and Wang Zhi Jian, The Ramsey Number $r(mC_4, nC_4)$ (in Chinese), *Journal of Shanghai Tiedao University*, **20** (1999) 66-70, 83.
- [LiWa2] Li Da Yong and Wang Zhi Jian, The Ramsey Numbers $r(mC_4, nC_5)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **45** (2003) 245-252.
- [-] Li Guiqing, see [SLLL, SLZL].
- [-] Li Jinwen, see [ZLLS].
- [-] Li Qiao, see [SLL, SLLL].
- [-] Li Wei, see [KLR].
- [Li1] Li Yusheng, Some Ramsey Numbers of Graphs with Bridge, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **25** (1997) 225-229.
- [Li2] Yusheng Li, Ramsey Numbers of a Cycle, *Taiwanese Journal of Mathematics*, **12** (2008) 1007-1013.
- [Li3] Yusheng Li, The Multi-Color Ramsey Number of an Odd Cycle, to appear in the *Journal of Graph Theory*, published online 2009.
- [LiLih] Yusheng Li and Ko-Wei Lih, Multi-Color Ramsey Numbers of Even Cycles, *European Journal of Combinatorics*, **30** (2009) 114-118.
- [LR1] Li Yusheng and C.C. Rousseau, On Book-Complete Graph Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **68** (1996) 36-44.
- [LR2] Li Yusheng and C.C. Rousseau, Fan-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **23** (1996) 413-420.
- [LR3] Li Yusheng and C.C. Rousseau, On the Ramsey Number $r(H + \bar{K}_n, K_n)$, *Discrete Mathematics*, **170** (1997) 265-267.
- [LR4] Li Yusheng and C.C. Rousseau, A Ramsey Goodness Result for Graphs with Many Pendant Edges, *Ars Combinatoria*, **49** (1998) 315-318.
- [LRS] Li Yusheng, C.C. Rousseau and L. Soltés, Ramsey Linear Families and Generalized Subdivided Graphs, *Discrete Mathematics*, **170** (1997) 269-275.
- [LiRZ1] Li Yusheng, C.C. Rousseau and Zang Wenan, Asymptotic Upper Bounds for Ramsey Functions, *Graphs and Combinatorics*, **17** (2001) 123-128.
- [LiRZ2] Li Yusheng, C.C. Rousseau and Zang Wenan, An Upper Bound on Ramsey Numbers, *Applied Mathematics Letters*, **17** (2004) 663-665.
- [LiShen] Yusheng Li and Jian Shen, Bounds for Ramsey Numbers of Complete Graphs Dropping an Edge, *European Journal of Combinatorics*, **29** (2008) 88-94.
- [LiTZ] Li Yusheng, Tang Xueqing and Zang Wenan, Ramsey Functions Involving $K_{m,n}$ with n Large, *Discrete Mathematics*, **300** (2005) 120-128.
- [LiZa1] Li Yusheng and Zang Wenan, Ramsey Numbers Involving Large Dense Graphs and Bipartite Turán Numbers, *Journal of Combinatorial Theory, Series B*, **87** (2003) 280-288.
- [LiZa2] Li Yusheng and Zang Wenan, The Independence Number of Graphs with a Forbidden Cycle and Ramsey Numbers, *Journal of Combinatorial Optimization*, **7** (2003) 353-359.
- [-] Li Yusheng, see also [BaiLi, BaLX, CLRZ, DoLL, GuLi, LinLi, ShiuLL, SunLi].

- [-] Li Zhenchong, see [LSL, LSWL].
- [-] Ko-Wei Lih, see [LiLih].
- [LinMac]* A. Lin and J. Mackey, Counting Triangles in Some Ramsey Graphs, *RIT Technical Report*, April (2009), <https://ritdml.rit.edu/dspace/handle/1850/8816>.
- [LinLi] Qizhong Lin and Yusheng Li, On Ramsey Numbers of Fans, *Discrete Applied Mathematics*, **157** (2009) 191-194.
- [-] Qizhong Lin, see also [DoLL].
- [-] Lin Xiaohui, see [SunYJLS, SunYLZ1, SunYLZ2].
- [Lind] B. Lindström, Undecided Ramsey-Numbers for Paths, *Discrete Mathematics*, **43** (1983) 111-112.
- [Ling] A.C.H. Ling, Some Applications of Combinatorial Designs to Extremal Graph Theory, *Ars Combinatoria*, **67** (2003) 221-229.
- [-] Andy Liu, see [AbbL].
- [-] Liu Linzhong, see [ZLLS].
- [-] Liu Shu Yan, see [SoBL].
- [-] Liu Xiangyang, see [GuSL].
- [-] Liu Yanwu, see [SoYL].
- [Loc] S.C. Locke, Bipartite Density and the Independence Ratio, *Journal of Graph Theory*, **10** (1986) 47-53.
- [-] S.C. Locke, see also [FrLo].
- [Lor] P.J. Lorimer, The Ramsey Numbers for Stripes and One Complete Graph, *Journal of Graph Theory*, **8** (1984) 177-184.
- [LorMu] P.J. Lorimer and P.R. Mullins, Ramsey Numbers for Quadrangles and Triangles, *Journal of Combinatorial Theory, Series B*, **23** (1977) 262-265.
- [LorSe] P.J. Lorimer and R.J. Segedin, Ramsey Numbers for Multiple Copies of Complete Graphs, *Journal of Graph Theory*, **2** (1978) 89-91.
- [LorSo] P.J. Lorimer and W. Solomon, The Ramsey Numbers for Stripes and Complete Graphs 1, *Discrete Mathematics*, **104** (1992) 91-97. Corrigendum in *Discrete Mathematics*, **131** (1994) 395.
- [-] P.J. Lorimer, see also [CL1, CL2].
- [Lortz] R. Lortz, A Note on the Ramsey Number of $K_{2,2}$ versus $K_{3,n}$, *Discrete Mathematics*, **306** (2006) 2976-2982.
- [LoM1] R. Lortz and I. Mengersen, On the Ramsey Numbers $r(K_{2,n-1}, K_{2,n})$ and $r(K_{2,n}, K_{2,n})$, *Utilitas Mathematica*, **61** (2002) 87-95.
- [LoM2] R. Lortz and I. Mengersen, Bounds on Ramsey Numbers of Certain Complete Bipartite Graphs, *Results in Mathematics*, **41** (2002) 140-149.
- [LoM3]* R. Lortz and I. Mengersen, Off-Diagonal and Asymptotic Results on the Ramsey Number $r(K_{2,m}, K_{2,n})$, *Journal of Graph Theory*, **43** (2003) 252-268.
- [LoM4]* R. Lortz and I. Mengersen, Further Ramsey Numbers for Small Complete Bipartite Graphs, *Ars Combinatoria*, **79** (2006) 195-203.
- [Łuc] T. Łuczak, $R(C_n, C_n, C_n) \leq (4+o(1))n$, *Journal of Combinatorial Theory, Series B*, **75** (1999) 174-187.
- [-] T. Łuczak, see also [FiŁu, HaŁP1+, HaŁP2+, HaŁT].
- [LSL]* Luo Haipeng, Su Wenlong and Li Zhenchong, The Properties of Self-Complementary Graphs and New Lower Bounds for Diagonal Ramsey Numbers, *Australasian Journal of Combinatorics*, **25** (2002) 103-116.
- [LSS1]* Luo Haipeng, Su Wenlong and Shen Yun-Qiu, New Lower Bounds of Ten Classical Ramsey Numbers, *Australasian Journal of Combinatorics*, **24** (2001) 81-90.

- [LSS2]* Luo Haipeng, Su Wenlong and Shen Yun-Qiu, New Lower Bounds for Two Multicolor Classical Ramsey Numbers, *Radovi Matematički*, **13** (2004) 15-21.
- [LSWL]* Luo Haipeng, Su Wenlong, Wu Kang and Li Zhenchong, Two New Lower Bounds for Ramsey Numbers $R(3, q)$ Based on the Parallel Algorithm (in Chinese), *Journal of Guangxi Academy of Sciences*, **19**, 4 (2003) 145-149.
- [-] Luo Haipeng, see also [SL, SLL, SLLL, SLZL, WSLX1, WSLX2].

M

- [Mac]* J. Mackey, Combinatorial Remedies, *Ph.D. Thesis*, Department of Mathematics, University of Hawaii, 1994.
- [] J. Mackey, see also [LinMac].
- [Mat]* R. Mathon, Lower Bounds for Ramsey Numbers and Association Schemes, *Journal of Combinatorial Theory, Series B*, **42** (1987) 122-127.
- [-] J.P. Mayberry, see [LayMa].
- [McS] C. McDiarmid and A. Steger, Tidier Examples for Lower Bounds on Diagonal Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **74** (1996) 147-152.
- [McK]** B.D. McKay, Australian National University, *personal communication* (2003). Graphs available at <http://cs.anu.edu.au/people/bdm/data/ramsey.html>.
- [MPR]** B.D. McKay, K. Piwakowski and S.P. Radziszowski, Ramsey Numbers for Triangles versus Almost-Complete Graphs, *Ars Combinatoria*, **73** (2004) 205-214.
- [MR1]** B.D. McKay and S.P. Radziszowski, The First Classical Ramsey Number for Hypergraphs is Computed, *Proceedings of the Second Annual ACM-SIAM Symposium on Discrete Algorithms, SODA'91*, San Francisco, (1991) 304-308.
- [MR2]* B.D. McKay and S.P. Radziszowski, A New Upper Bound for the Ramsey Number $R(5, 5)$, *Australian Journal of Combinatorics*, **5** (1992) 13-20.
- [MR3]** B.D. McKay and S.P. Radziszowski, Linear Programming in Some Ramsey Problems, *Journal of Combinatorial Theory, Series B*, **61** (1994) 125-132.
- [MR4]** B.D. McKay and S.P. Radziszowski, $R(4, 5) = 25$, *Journal of Graph Theory*, **19** (1995) 309-322.
- [MR5]** B.D. McKay and S.P. Radziszowski, Subgraph Counting Identities and Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **69** (1997) 193-209.
- [MZ]** B.D. McKay and Zhang Ke Min, The Value of the Ramsey Number $R(3, 8)$, *Journal of Graph Theory*, **16** (1992) 99-105.
- [-] B.D. McKay, see also [FM].
- [McN]** J. McNamara, SUNY Brockport, *personal communication* (1995).
- [McR]** J. McNamara and S.P. Radziszowski, The Ramsey Numbers $R(K_4 - e, K_6 - e)$ and $R(K_4 - e, K_7 - e)$, *Congressus Numerantium*, **81** (1991) 89-96.
- [MO] I. Mengersen and J. Oeckermann, Matching-Star Ramsey Sets, *Discrete Applied Mathematics*, **95** (1999) 417-424.
- [-] I. Mengersen, see also [AKM, CEHMS, EHM1, EHM2, HoMe, HaMe1, HaMe2, HaMe3, HaMe4, KM1, KM2, LoM1, LoM2, LoM3, LoM4].
- [-] Zhengke Miao, see [ChenCMN].
- [-] M. Miller, see [BSNM].
- [MiSa] H. Mizuno and I. Sato, Ramsey Numbers for Unions of Some Cycles, *Discrete Mathematics*, **69** (1988) 283-294.
- [MoCa] E.L. Monte Carmelo, Configurations in Projective Planes and Quadrilateral-Star Ramsey Numbers, *Discrete Mathematics*, **308** (2008) 3986-3991.

- [-] E.L. Monte Carmelo, see also [GoMC].
- [-] D. Mubayi, see [AFM, LaMu].
- [-] P.R. Mullins, see [LorMu].

N

- [-] S.M. Nababan, see [BSNM].
- [NaORS] B. Nagle, S. Olsen, V. Rödl and M. Schacht, On the Ramsey Number of Sparse 3-Graphs, *Graphs and Combinatorics*, **24** (2008) 205-228.
- [Neš] J. Nešetřil, Ramsey Theory, chapter 25 in *Handbook of Combinatorics*, ed. R.L. Graham, M. Grötschel and L. Lovász, The MIT-Press, Vol. II, 1996, 1331-1403.
- [-] J. Nešetřil, see also [GrNe].
- [-] C.T. Ng, see [ChenCMN, ChenCN, CheCZN].
- [Nik] V. Nikiforov, The Cycle-Complete Graph Ramsey Numbers, *Combinatorics, Probability and Computing*, **14** (2005) 349-370.
- [NiRo1] V. Nikiforov and C.C. Rousseau, Large Generalized Books Are p -Good, *Journal of Combinatorial Theory, Series B*, **92** (2004) 85-97.
- [NiRo2] V. Nikiforov and C.C. Rousseau, Book Ramsey Numbers I, *Random Structures and Algorithms*, **27** (2005) 379-400.
- [NiRo3] V. Nikiforov and C.C. Rousseau, A Note on Ramsey Numbers for Books, *Journal of Graph Theory*, **49** (2005) 168-176.
- [NiRo4] V. Nikiforov and C.C. Rousseau, Ramsey Goodness and Beyond, to appear in *Combinatorica*.
- [NiRS] V. Nikiforov, C.C. Rousseau and R.H. Schelp, Book Ramsey Numbers and Quasi-Randomness, *Combinatorics, Probability and Computing*, **14** (2005) 851-860.
- [-] A. Nowik, see [DzNS].

O

- [-] J. Oeckermann, see [MO].
- [-] S. Olsen, see [NaORS].
- [-] D. Osthus, see [CooFKO, KüCFO].

P

- [-] Linqiang Pan, see [ShaXBP, ShaXP, ShaXSP].
- [Par1] T.D. Parsons, The Ramsey Numbers $r(P_m, K_n)$, *Discrete Mathematics*, **6** (1973) 159-162.
- [Par2] T.D. Parsons, Path-Star Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **17** (1974) 51-58.
- [Par3] T.D. Parsons, Ramsey Graphs and Block Designs, I, *Transactions of the American Mathematical Society*, **209** (1975) 33-44.
- [Par4] T.D. Parsons, Ramsey Graphs and Block Designs, *Journal of Combinatorial Theory, Series A*, **20** (1976) 12-19.
- [Par5] T.D. Parsons, Graphs from Projective Planes, *Aequationes Mathematicae*, **14** (1976) 167-189.
- [Par6] T.D. Parsons, Ramsey Graph Theory, in *Selected Topics in Graph Theory*, (L.W. Beineke and R.J. Wilson eds.), Academic Press, (1978) 361-384.
- [-] T.D. Parsons, see also [FLPS].

- [-] Y. Peng, see [HaLP1+, HaLP2+].
- [-] O. Pikhurko, see [BePi].
- [Piw1]* K. Piwakowski, Applying Tabu Search to Determine New Ramsey Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R6, **3** (1996), 4 pages.
- [Piw2]** K. Piwakowski, A New Upper Bound for $R_3(K_4-e)$, *Congressus Numerantium*, **128** (1997) 135-141.
- [PR1]** K. Piwakowski and S.P. Radziszowski, $30 \leq R(3,3,4) \leq 31$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **27** (1998) 135-141.
- [PR2]** K. Piwakowski and S.P. Radziszowski, Towards the Exact Value of the Ramsey Number $R(3,3,4)$, *Congressus Numerantium*, **148** (2001) 161-167.
- [-] K. Piwakowski, see also [MPR, DzKP].
- [PoRRS] J. Polcyn, V. Rödl, A. Ruciński and E. Szemerédi, Short Paths in Quasi-Random Triple Systems with Sparse Underlying Graphs, *Journal of Combinatorial Theory, Series B*, **96** (2006) 584-607.
- [-] A.D. Polimeni, see [CGP, CRSPS].
- [-] L.M. Pretorius, see [SwPr].
- [-] P. Pudlák, see [AlPu, CPR].

Q

- [-] Qian Xinjin, see [SoGQ].

R

- [Ra1]** S.P. Radziszowski, The Ramsey Numbers $R(K_3, K_8-e)$ and $R(K_3, K_9-e)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **8** (1990) 137-145.
- [Ra2] S.P. Radziszowski, Small Ramsey Numbers, *Technical Report RIT-TR-93-009*, Department of Computer Science, Rochester Institute of Technology (1993).
- [Ra3]** S.P. Radziszowski, On the Ramsey Number $R(K_5-e, K_5-e)$, *Ars Combinatoria*, **36** (1993) 225-232.
- [RaJi] S.P. Radziszowski and Jin Xia, Paths, Cycles and Wheels in Graphs without Antitriangles, *Australasian Journal of Combinatorics*, **9** (1994) 221-232.
- [RK1]* S.P. Radziszowski and D.L. Kreher, Search Algorithm for Ramsey Graphs by Union of Group Orbits, *Journal of Graph Theory*, **12** (1988) 59-72.
- [RK2]** S.P. Radziszowski and D.L. Kreher, Upper Bounds for Some Ramsey Numbers $R(3, k)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 207-212.
- [RK3]** S.P. Radziszowski and D.L. Kreher, On $R(3, k)$ Ramsey Graphs: Theoretical and Computational Results, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 37-52.
- [RK4] S.P. Radziszowski and D.L. Kreher, Minimum Triangle-Free Graphs, *Ars Combinatoria*, **31** (1991) 65-92.
- [RT]* S.P. Radziszowski and Kung-Kuen Tse, A Computational Approach for the Ramsey Numbers $R(C_4, K_n)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **42** (2002) 195-207.
- [RST]* S.P. Radziszowski, J. Stinehour and Kung-Kuen Tse, Computation of the Ramsey Number $R(W_5, K_5)$, *Bulletin of the Institute of Combinatorics and its Applications*, **47** (2006) 53-57.
- [-] S.P. Radziszowski, see also [BaRT, BLR, FKR, KLR, MPR, MR1, MR2, MR3, MR4, MR5, McR, PR1, PR2, XuR1, XuR2, XSR1, XSR2, XXER, XXR].
- [Ram] F.P. Ramsey, On a Problem of Formal Logic, *Proceedings of the London Mathematical Society*, **30** (1930) 264-286.

- [RaHo]** Ramsey@Home, A distributed computing project searching for lower bounds for Ramsey numbers, <http://www.ramseyathome.com/ramsey> (2009).
- [Rao]* S. Rao, Applying a Genetic Algorithm to Improve the Lower Bounds of Multi-Color Ramsey Numbers, *MS thesis*, Department of Computer Science, Rochester Institute of Technology, 1997.
- [-] G. Resta, see [CPR].
- [-] S.W. Reyner, see [BR].
- [-] D.F. Reynolds, see [ExRe].
- [Rob1] F.S. Roberts, *Applied Combinatorics*, Prentice-Hall, Englewood Cliffs, 1984.
- [-] J.A. Roberts, see [BuRo1, BuRo2].
- [-] S. Roberts, see [GR].
- [Rob2]* A. Robertson, New Lower Bounds for Some Multicolored Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R12, **6** (1999), 6 pages.
- [Rob3]* A. Robertson, Difference Ramsey Numbers and Issai Numbers, *Advances in Applied Mathematics*, **25** (2000) 153-162.
- [Rob4] A. Robertson, New Lower Bounds Formulas for Multicolored Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R13, **9** (2002), 6 pages.
- [-] Y. Roditty, see [KrRod].
- [RöTh] V. Rödl and R. Thomas, Arrangeability and Clique Subdivisions, in *The Mathematics of Paul Erdős II*, 236-239, Algorithms and Combinatorics **14**, Springer, Berlin, 1997.
- [-] V. Rödl, see also [AIRö, CRST, DLR, GrRö, GRR1, GRR2, HaŁP1+, HaŁP2+, KoRö1, KoRö2, KoRö3, NaORS, PoRRS].
- [Ros1] V. Rosta, On a Ramsey Type Problem of J.A. Bondy and P. Erdős, I & II, *Journal of Combinatorial Theory, Series B*, **15** (1973) 94-120.
- [Ros2] V. Rosta, Ramsey Theory Applications, Dynamic Survey in *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/Surveys>, #DS13, (2004), 43 pages.
- [-] V. Rosta, see also [BuRo3, KáRos].
- [-] B.L. Rothschild, see [GRS].
- [Rou] C.C. Rousseau, *personal communication*, (2006).
- [RoJa1] C.C. Rousseau and C.J. Jayawardene, The Ramsey Number for a Quadrilateral vs. a Complete Graph on Six Vertices, *Congressus Numerantium*, **123** (1997) 97-108.
- [RoJa2] C.C. Rousseau and C.J. Jayawardene, Harary's Problem for $K_{2,k}$, *unpublished manuscript*, (1999).
- [RS1] C.C. Rousseau and J. Sheehan, On Ramsey Numbers for Books, *Journal of Graph Theory*, **2** (1978) 77-87.
- [RS2] C.C. Rousseau and J. Sheehan, A Class of Ramsey Problems Involving Trees, *Journal of the London Mathematical Society (2)*, **18** (1978) 392-396.
- [-] C.C. Rousseau, see also [BJYHRZ, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, CLRZ, CRSPS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9, FRS1, FRS2, FRS3, FRS4, FRS5, FRS6, FRS7, FRS8, FSR, JR1, JR2, JR3, JR4, JR5, LR1, LR2, LR3, LR4, LRS, LiRZ1, LiRZ2, NiRo1, NiRo2, NiRo3, NiRo4, NiRS].
- [-] C. Rowan, see [KerRo].
- [-] P. Rowlinson, see [YR1, YR2, YR3].
- [-] A. Ruciński, see [GRR1, GRR2, HaŁP1+, HaŁP2+, PoRRS].
- [-] M. Ruszinkó, see [GyRSS].

S

- [SaBr1] A.N.M. Salman and H.J. Broersma, The Ramsey Numbers of Paths versus Kipases, *Electronic Notes in Discrete Mathematics*, **17** (2004) 251-255.
- [SaBr2] A.N.M. Salman and H.J. Broersma, Paths-Fan Ramsey Numbers, *Discrete Applied Mathematics*, **154** (2006) 1429-1436.
- [SaBr3] A.N.M. Salman and H.J. Broersma, The Ramsey Numbers for Paths versus Wheels, *Discrete Mathematics*, **307** (2007) 975-982.
- [SaBr4] A.N.M. Salman and H.J. Broersma, Path-Kipas Ramsey Numbers, *Discrete Applied Mathematics*, **155** (2007) 1878-1884.
- [San] A. Sánchez-Flores, An Improved Bound for Ramsey Number $N(3,3,3,3;2)$, *Discrete Mathematics*, **140** (1995) 281-286.
- [-] G.N. Sárközy, see [GyRSS, GySS].
- [-] I. Sato, see [MiSa].
- [-] M. Schacht, see [NaORS].
- [Scha] M. Schaefer, Graph Ramsey Theory and the Polynomial Hierarchy, *Journal of Computer and System Sciences*, **62** (2001) 290-322.
- [-] R.H. Schelp, see [BaLS, BaSS, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, BEFS, ChenS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9, FLPS, FRS1, FRS2, FRS3, FRS4, FRS5, FS1, FS2, FS3, FS4, FSR, FSS1, NiRS].
- [-] J. Schönheim, see [BS].
- [SchSch1]* A. Schelten and I. Schiermeyer, Ramsey Numbers $r(K_3, G)$ for Connected Graphs G of Order Seven, *Discrete Applied Mathematics*, **79** (1997) 189-200.
- [SchSch2] A. Schelten and I. Schiermeyer, Ramsey Numbers $r(K_3, G)$ for $G \cong K_7 - 2P_2$ and $G \cong K_7 - 3P_2$, *Discrete Mathematics*, **191** (1998) 191-196.
- [-] A. Schelten, see also [FSS2].
- [Schi1] I. Schiermeyer, All Cycle-Complete Graph Ramsey Numbers $r(C_m, K_6)$, *Journal of Graph Theory*, **44** (2003) 251-260.
- [Schi2] I. Schiermeyer, The Cycle-Complete Graph Ramsey Number $r(C_5, K_7)$, *Discussiones Mathematicae Graph Theory*, **25** (2005) 129-139.
- [-] I. Schiermeyer, see also [FSS2, SchSch1, SchSch2].
- [Schu] C. -U. Schulte, Ramsey-Zahlen für Bäume und Kreise, *Ph.D. thesis*, Heinrich-Heine-Universität Düsseldorf, (1992).
- [-] S. Schuster, see [ChaS].
- [-] A. Schwenk, see [ChvS].
- [Scob] M.W. Scobee, On the Ramsey Number $R(m_1P_3, m_2P_3, m_3P_3)$ and Related Results, ..., *MA thesis*, University of Louisville (1993).
- [-] R.J. Segegin, see [LorSe].
- [Shao]* Zehui Shao, *personal communication* (2008).
- [ShaXBP]* Zehui Shao, Jin Xu, Qiquan Bao and Linqiang Pan, Computation of Some Generalized Ramsey Numbers, to appear in the *Journal of Combinatorial Mathematics and Combinatorial Computing*.
- [ShaXP]* Zehui Shao, Jin Xu and Linqiang Pan, Lower Bounds on Ramsey Numbers $R(6,8)$, $R(7,9)$ and $R(8,17)$, to appear in *Ars Combinatoria*.
- [ShaXSP]* Zehui Shao, Xiaodong Xu, Xiaolong Shi and Linqiang Pan, Some Three-Color Ramsey Numbers $R(P_4, P_5, C_k)$ and $R(P_4, P_6, C_k)$, *European Journal of Combinatorics*, **30** (2009) 396-403.
- [-] Zehui Shao, see also [XSR1, XSR2].

- [Shas] A. Shastri, Lower Bounds for Bi-Colored Quaternary Ramsey Numbers, *Discrete Mathematics*, **84** (1990) 213-216.
- [She1]* J.B. Shearer, Lower Bounds for Small Diagonal Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **42** (1986) 302-304.
- [She2] J.B. Shearer, A Note on the Independence Number of Triangle-free Graphs II, *Journal of Combinatorial Theory, Series B*, **53** (1991) 300-307.
- [She3]* J.B. Shearer, Independence Numbers of Paley Graphs (data for primes $1 \pmod{4}$ up to 7000), <http://www.research.ibm.com/people/s/shearer/indpal.html> (1996).
- [-] J. Sheehan, see [CRSPS, CEHMS, FRS6, FRS7, FRS8, RS1, RS2].
- [-] Jian Shen, see [LiShen].
- [-] Shen Yun-Qiu, see [LSS1, LSS2].
- [-] Sheng Wancheng, see [HWSYZH].
- [-] Shi Lei, see [SunYJLS].
- [Shi1] Shi Ling Sheng, Cube Ramsey Numbers Are Polynomial, *Random Structures & Algorithms*, **19** (2001) 99--101.
- [Shi2] Shi Ling Sheng, Upper Bounds for Ramsey Numbers, *Discrete Mathematics*, **270** (2003) 251-265.
- [Shi3] Shi Ling Sheng, Linear Ramsey Numbers of Sparse Graphs, *Journal of Graph Theory*, **50** (2005) 175-185.
- [Shi4] Shi Lingsheng, The Tail Is Cut for Ramsey Numbers of Cubes, *Discrete Mathematics*, **307** (2007) 290-292.
- [ShZ1] Shi Ling Sheng and Zhang Ke Min, An Upper Bound Formula for Ramsey Numbers, *manuscript*, (2001).
- [ShZ2] Shi Ling Sheng and Zhang Ke Min, A Sequence of Formulas for Ramsey Numbers, *manuscript*, (2001).
- [-] Xiaolong Shi, see [ShaXSP].
- [ShiuLL] Shiu Wai Chee, Peter Che Bor Lam and Li Yusheng, On Some Three-Color Ramsey Numbers, *Graphs and Combinatorics*, **19** (2003) 249-258.
- [Sid1] A.F. Sidorenko, On Turán Numbers $T(n, 5, 4)$ and Number of Monochromatic 4-cliques in 2-colored 3-graphs (in Russian), *Voprosy Kibernetiki*, **64** (1980) 117-124.
- [Sid2] A.F. Sidorenko, An Upper Bound on the Ramsey Number $R(K_3, G)$ Depending Only on the Size of the Graph G , *Journal of Graph Theory*, **15** (1991) 15-17.
- [Sid3] A.F. Sidorenko, The Ramsey Number of an N -Edge Graph versus Triangle Is at Most $2N + 1$, *Journal of Combinatorial Theory, Series B*, **58** (1993) 185-196.
- [-] M. Simonovits, see [BaSS, FSS1, FS, HaLP1+, KoSS].
- [SiqSk] F. Siqueira Benevides and J. Skokan, The 3-Colored Ramsey Number of Even Cycles, *Journal of Combinatorial Theory, Series B*, **99** (2009) 690-708.
- [-] J. Skokan, see [HaLP1+, HaLP2+, KoSS, SiqSk].
- [-] M.J. Smuga-Otto, see [AbbS].
- [Sob] A. Sobczyk, Euclidian Simplices and the Ramsey Number $R(4, 4; 3)$, *Technical Report #10, Clemson University* (1967).
- [Soi] A. Soifer, *The Mathematical Coloring Book*, Springer, 2009.
- [-] W. Solomon, see [LorSo].
- [-] L. Soltés, see [LRS].
- [Song1] Song En Min, Study of Some Ramsey Numbers (in Chinese), a note (announcement of results without proofs), *Mathematica Applicata*, **4(2)** (1991) 6.

- [Song2] Song En Min, New Lower Bound Formulas for the Ramsey Numbers $N(k, k, \dots, k; 2)$ (in Chinese), *Mathematica Applicata*, **6** (1993) suppl., 113-116.
- [Song3] Song En Min, An Investigation of Properties of Ramsey Numbers (in Chinese), *Mathematica Applicata*, **7** (1994) 216-221.
- [Song4] Song En Min, Properties and New Lower Bounds of the Ramsey Numbers $R(p, q; 4)$ (in Chinese), *Journal of Huazhong University of Science and Technology*, **23** (1995) suppl. II, 1-4.
- [SoYL] Song En Min, Ye Weiguo and Liu Yanwu, New Lower Bounds for Ramsey Number $R(p, q; 4)$, *Discrete Mathematics*, **145** (1995) 343-346.
- [-] Song En Min, see also [HuSo, ZLLS].
- [Song5] Song Hongxue, Asymptotic Upper Bounds for Wheel-Complete Graph Ramsey Numbers, *Journal of Southeast University* (English Edition), ISSN 1003-7985, **20** (2004) 126-129.
- [SoBL] Song Hong Xue, Bai Lu Feng and Liu Shu Yan, Asymptotic Upper Bounds for the Wheel-Complete Graph Ramsey Numbers (in Chinese), *Acta Mathematica Scientia, Series A*, ISSN 1003-3998, **26** (2006) 741-746.
- [SoGQ] Song Hongxue, Gu Hua and Qian Xinjin, On the Ramsey Number of K_3 versus $K_2 + T_n$ (in Chinese), *Journal of Liaoning Normal University, Natural Science Edition*, ISSN 1000-1735, **27** (2004) 142-145.
- [-] Song Hongxue, see also [GuSL].
- [Spe1] J.H. Spencer, Ramsey's Theorem - A New Lower Bound, *Journal of Combinatorial Theory, Series A*, **18** (1975) 108-115.
- [Spe2] J.H. Spencer, Asymptotic Lower Bounds for Ramsey Functions, *Discrete Mathematics*, **20** (1977/1978) 69-76.
- [-] J.H. Spencer, see also [BES, GRS].
- [Spe3]* T. Spencer, University of Nebraska at Omaha, *personal communication* (1993), and, Upper Bounds for Ramsey Numbers via Linear Programming, *manuscript*, (1994).
- [-] A.K. Srivastava, see [GauST].
- [Stahl] S. Stahl, On the Ramsey Number $R(F, K_m)$ where F is a Forest, *Canadian Journal of Mathematics*, **27** (1975) 585-589.
- [-] R.G. Stanton, see [KaSt].
- [Stat] W. Staton, Some Ramsey-type Numbers and the Independence Ratio, *Transactions of the American Mathematical Society*, **256** (1979) 353-370.
- [-] A. Steger, see [McS].
- [-] J. Stinehour, see [RST].
- [Stev] S. Stevens, Ramsey Numbers for Stars versus Complete Multipartite Graphs, *Congressus Numerantium*, **73** (1990) 63-71.
- [-] M.J. Stewart, see [CRSPS].
- [Stone] J.C. Stone, Utilizing a Cancellation Algorithm to Improve the Bounds of $R(5, 5)$, (1996), <http://oas.okstate.edu/ojas/jstone.htm>. This paper claims incorrectly that $R(5, 5) = 50$.
- [SL]* Su Wenlong and Luo Haipeng, Prime Order Cyclic Graphs and New Lower Bounds for Three Classical Ramsey Numbers $R(4, n)$ (in Chinese), *Journal of Mathematical Study*, **31**, 4 (1998) 442-446.
- [SLL]* Su Wenlong, Luo Haipeng and Li Qiao, New Lower Bounds of Classical Ramsey Numbers $R(4, 12)$, $R(5, 11)$ and $R(5, 12)$, *Chinese Science Bulletin*, **43**, 6 (1998) 528.
- [SLLL]* Su Wenlong, Luo Haipeng, Li Guiqing and Li Qiao, Lower Bounds of Ramsey Numbers Based on Cubic Residues, *Discrete Mathematics*, **250** (2002) 197-209.
- [SLZL]* Su Wenlong, Luo Haipeng, Zhang Zhengyou and Li Guiqing, New Lower Bounds of Fifteen Classical Ramsey Numbers, *Australasian Journal of Combinatorics*, **19** (1999) 91-99.

- [–] Su Wenlong, see also [LSL, LSS1, LSS2, LSWL, WuCXS, WSLX1, WSLX2].
- [Sud1] B. Sudakov, A Note on Odd Cycle-Complete Graph Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #N1, **9** (2002), 4 pages.
- [Sud2] B. Sudakov, Large K_r -free Subgraphs in K_s -free Graphs and Some Other Ramsey-Type Problems, *Random Structures and Algorithms*, **26** (2005) 253-265.
- [Sud3] B. Sudakov, Ramsey Numbers and the Size of Graphs, *SIAM Journal on Discrete Mathematics*, **21** (2007) 980--986.
- [–] B. Sudakov, see also [AIKS, ConFS1, ConFS2, ConFS3, FoxSu1, FoxSu2, KoSu].
- [–] A. Sudan, see [GGS].
- [Sun]* Sun Yongqi, Research on Ramsey Numbers of Some Graphs (in Chinese), *Ph. D. thesis*, Dalian University of Technology, China, July 2006.
- [SunYJLS] Sun Yongqi, Yang Yuansheng, Jiang Baoqi, Lin Xiaohui and Shi Lei, On Multicolor Ramsey Numbers for Even Cycles in Graphs, *Ars Combinatoria*, **84** (2007) 333-343.
- [SunYLZ1]* Sun Yongqi, Yang Yuansheng, Lin Xiaohui and Zheng Wenping, The Value of the Ramsey Number $R_4(C_4)$, *Utilitas Mathematica*, **73** (2007) 33-44.
- [SunYLZ2]* Sun Yongqi, Yang Yuansheng, Lin Xiaohui and Zheng Wenping, On the Three Color Ramsey Numbers $R(C_m, C_4, C_4)$, *Ars Combinatoria*, **84** (2007) 3-11.
- [SunYW]* Sun Yongqi, Yang Yuansheng and Wang Zhihai, The Value of the Ramsey Number $R_5(C_6)$, *Utilitas Mathematica*, **76** (2008) 25-31.
- [SunYWLX]* Sun Yongqi, Yang Yuansheng, Wang Wei, Li Bingxi and Xu Feng, Study of Three Color Ramsey numbers $R(C_{m_1}, C_{m_2}, C_{m_3})$ (in Chinese), *Journal of Dalian University of Technology*, ISSN 1000-8608, **46** (2006) 428-433.
- [SunYXL] Sun Yongqi, Yang Yuansheng, Xu Feng and Li Bingxi, New Lower Bounds on the Multicolor Ramsey Numbers $R_r(C_{2m})$, *Graphs and Combinatorics*, **22** (2006) 283-288.
- [SunLi] Sun Yuqin and Li Yusheng, On an Upper Bound of Ramsey Number $r_k(K_{m,n})$ with Large n , *Heilongjiang Daxue Ziran Kexue Xuebao*, ISSN 1001-7011, **23** (2006) 668-670.
- [Sur] Surahmat, Cycle-Wheel Ramsey Numbers. Some results, open problems and conjectures. *Math Track*, ISSN 1817-3462, 1818-5495, **2** (2006) 56-64.
- [SuBa1] Surahmat and E.T. Baskoro, On the Ramsey Number of a Path or a Star versus W_4 or W_5 , *Proceedings of the 12-th Australasian Workshop on Combinatorial Algorithms*, Bandung, Indonesia, July 14-17 (2001) 174-179.
- [SuBa2] Surahmat and E.T. Baskoro, The Ramsey Number of Linear Forest versus Wheel, paper presented at the *13-th Australasian Workshop on Combinatorial Algorithms*, Fraser Island, Queensland, Australia, July 7-10, 2002.
- [SuBB1] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Star-like Trees versus Large Odd Wheels, *Technical Report #1621*, Faculty of Mathematical Sciences, University of Twente, The Netherlands, (2002).
- [SuBB2] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Cycles versus Small Wheels, *Integers: Electronic Journal of Combinatorial Number Theory*, <http://www.integers-ejcnt.org/vol4.html>, #A10, **4** (2004), 9 pages.
- [SuBB3] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Fans versus K_4 , *Bulletin of the Institute of Combinatorics and its Applications*, **43** (2005) 96-102.
- [SuBB4] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Star and Large Star-Like Trees versus Odd Wheels, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **65** (2008) 153-162.
- [SuBT1] Surahmat, E.T. Baskoro and I. Tomescu, The Ramsey Numbers of Large Cycles versus Wheels, *Discrete Mathematics*, **306** (2006), 3334-3337.

- [SuBT2] Surahmat, E.T. Baskoro and I. Tomescu, The Ramsey Numbers of Large Cycles versus Odd Wheels, *Graphs and Combinatorics*, **24** (2008), 53-58.
- [SuBTB] Surahmat, E.T. Baskoro, I. Tomescu and H.J. Broersma, On Ramsey Numbers of Cycles with Respect to Generalized Even Wheels, *manuscript*, (2006).
- [SuBUB] Surahmat, E.T. Baskoro, S. Uttunggadewa and H.J. Broersma, An Upper Bound for the Ramsey Number of a Cycle of Length Four versus Wheels, in *LNCS 3330*, Springer, Berlin (2005) 181-184.
- [-] Surahmat, see also [AliSur, BaSu, BSNM].
- [SwPr] C.J. Swanepoel and L.M. Pretorius, Upper Bounds for a Ramsey Theorem for Trees, *Graphs and Combinatorics*, **10** (1994) 377-382.
- [-] M.M. Sweet, see [FreSw].
- [-] E. Szemerédi, see [AKS, CRST, GyRSS, GySS, PoRRS].
- [-] P. Szuca, see [DzNS].

T

- [-] Tang Xueqing, see [LiTZ].
- [-] R. Thomas, see [RöTh].
- [Tho] A. Thomason, An Upper Bound for Some Ramsey Numbers, *Journal of Graph Theory*, **12** (1988) 509-517.
- [-] P.W. Tingley, see [HaLT].
- [-] I. Tomescu, see [AliBT, SuBT1, SuBT2, SuBTB].
- [-] C.A. Tovey, see [CET].
- [-] A. Tripathi, see [GauST].
- [Tr] Trivial results.
- [-] W.T. Trotter Jr., see [CRST].
- [Tse1]* Kung-Kuen Tse, On the Ramsey Number of the Quadrilateral versus the Book and the Wheel, *Australasian Journal of Combinatorics*, **27** (2003) 163-167.
- [Tse2]* Kung-Kuen Tse, A Note on the Ramsey Numbers $R(C_4, B_n)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **58** (2006) 97-100.
- [Tse3]* Kung-Kuen Tse, A Note on Some Ramsey Numbers $R(C_p, C_q, C_r)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **62** (2007) 189-192.
- [-] Kung-Kuen Tse, see also [BaRT, RST, RT].
- [-] Z. Tuza, see [GyTu].

U

- [-] S. Uttunggadewa, see [SuBUB].

V

- [-] L. Volkmann, see [GuoV].

W

- [Walk] K. Walker, Dichromatic Graphs and Ramsey Numbers, *Journal of Combinatorial Theory*, **5** (1968) 238-243.

- [Wall] W.D. Wallis, On a Ramsey Number for Paths, *Journal of Combinatorics, Information & System Sciences*, **6** (1981) 295-296.
- [Wan] Wan Honghui, Upper Bounds for Ramsey Numbers $R(3, 3, \dots, 3)$ and Schur Numbers, *Journal of Graph Theory*, **26** (1997) 119-122.
- [-] Wang Gongben, see [WW, WWY1, WWY2].
- [WW]* Wang Qingxian and Wang Gongben, New Lower Bounds of Ramsey Numbers $r(3, q)$ (in Chinese), *Acta Scientiarum Naturalium, Universitatis Pekinensis*, **25** (1989) 117-121.
- [WWY1]* Wang Qingxian, Wang Gongben and Yan Shuda, A Search Algorithm And New Lower Bounds for Ramsey Numbers $r(3, q)$, *manuscript*, (1994).
- [WWY2]* Wang Qingxian, Wang Gongben and Yan Shuda, The Ramsey Numbers $R(K_3, K_q - e)$ (in Chinese), *Beijing Daxue Xuebao Ziran Kexue Ban*, **34** (1998) 15-20.
- [-] Wang Wei, see [SunYWLX, SunYXL].
- [-] Wang Yuandi, see [HWSYZH].
- [-] Wang Zhihai, see [SunYW].
- [-] Wang Zhi Jian, see [LiWa1, LiWa2].
- [West] D. West, *Introduction to Graph Theory*, second edition, Prentice Hall, 2001.
- [Wh] E.G. Whitehead, The Ramsey Number $N(3, 3, 3, 3; 2)$, *Discrete Mathematics*, **4** (1973) 389-396.
- [-] E.R. Williams, see [AbbW].
- [-] R.M. Wilson, see [FraWi].
- [-] A. Woldar, see [LaWo1, LaWo2].
- [WuCXS]* Wu Kang, Chen Hong, Xu Xiaodong and Su Wenlong, New Lower Bound for Nine Classical Ramsey Numbers $R(3, t)$ (in Chinese), *submitted*.
- [WSLX1]* Wu Kang, Su Wenlong, Luo Haipeng and Xu Xiaodong, New Lower Bound for Seven Classical Ramsey Numbers $R(3, q)$, *Applied Mathematics Letters*, **22** (2009) 365-368.
- [WSLX2]* Wu Kang, Su Wenlong, Luo Haipeng and Xu Xiaodong, A Generalization of Generalized Paley Graphs and New Lower Bounds for $R(3, q)$, *manuscript*, (2009).
- [-] Wu Kang, see also [LSWL].

X

- [XieZ]* Xie Jiguo and Zhang Xiaoxian, A New Lower Bound for Ramsey Number $r(3, 13)$ (in Chinese), *Journal of Lanzhou Railway Institute*, **12** (1993) 87-89.
- [-] Xie Zheng, see [XX1, XX2, XXER, XXR].
- [-] Jin Xu, see [ShaXBP, ShaXP].
- [-] Xu Feng, see [SunYWLX, SunYXL].
- [-] Ran Xu, see [ChenCX].
- [Xu] Xu Xiaodong, *personal communication*, (2004).
- [XuR1] Xiaodong Xu and S.P. Radziszowski, An Improvement to MATHON's Cyclotomic Ramsey Colorings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #N1, **16(1)** (2009), 5 pages.
- [XuR2] Xiaodong Xu and S.P. Radziszowski, $28 \leq R(C_4, C_4, C_3, C_3) \leq 36$, to appear in *Utilitas Mathematica*.
- [XSR1]* Xiaodong Xu, Zehui Shao and S.P. Radziszowski, Bounds on Some Ramsey Numbers Involving Quadrilateral, *Ars Combinatoria*, **90** (2009) 337-344.
- [XSR2]* Xiaodong Xu, Zehui Shao and S.P. Radziszowski, More Constructive Lower Bounds on Classical Ramsey Numbers, *in preparation*, (2009).

- [XX1]* Xu Xiaodong and Xie Zheng, A Constructive Approach for the Lower Bounds on the Ramsey Numbers $r(k, l)$, *manuscript*, (2002).
- [XX2] Xu Xiaodong and Xie Zheng, A Constructive Approach for the Lower Bounds on Multicolor Ramsey Numbers, *manuscript*, (2002).
- [XXER]* Xu Xiaodong, Xie Zheng, G. Exoo and S.P. Radziszowski, Constructive Lower Bounds on Classical Multicolor Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R35, **11** (2004), 24 pages.
- [XXR] Xu Xiaodong, Xie Zheng and S.P. Radziszowski, A Constructive Approach for the Lower Bounds on the Ramsey Numbers $R(s, t)$, *Journal of Graph Theory*, **47** (2004) 231-239.
- [-] Xu Xiaodong, see also [ShaXSP, WuCXS, WSLX1, WSLX2].
- [-] Xu Zhiqiang, see [BaLX].

Y

- [-] J. Yackel, see [GY].
- [-] Yan Shuda, see [WWY1, WWY2].
- [Yang] Yang Jian Sheng, results which can be obtained by the methods in [HWSYZH], *personal communication*, (2005).
- [YHZ1] Yang Jian Sheng, Huang Yi Ru and Zhang Ke Min, The Value of the Ramsey Number $R(C_n, K_4)$ is $3(n-1)+1$ ($n \geq 4$), *Australasian Journal of Combinatorics*, **20** (1999) 205-206.
- [YHZ2] Yang Jian Sheng, Huang Yi Ru and Zhang Ke Min, $R(C_6, K_5) = 21$ and $R(C_7, K_5) = 25$, *European Journal of Combinatorics*, **22** (2001) 561-567.
- [-] Yang Jian Sheng, see also [BJYHRZ, HWSYZH, HYZ].
- [YY]** Yang Yuansheng, On the Third Ramsey Numbers of Graphs with Six Edges, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **17** (1995) 199-208.
- [YH]* Yang Yuansheng and G.R.T. Hendry, The Ramsey Number $r(K_1 + C_4, K_5 - e)$, *Journal of Graph Theory*, **19** (1995) 13-15.
- [YR1]** Yang Yuansheng and P. Rowlinson, On the Third Ramsey Numbers of Graphs with Five Edges, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **11** (1992) 213-222.
- [YR2]* Yang Yuansheng and P. Rowlinson, On Graphs without 6-Cycles and Related Ramsey Numbers, *Utilitas Mathematica*, **44** (1993) 192-196.
- [YR3]* Yang Yuansheng and P. Rowlinson, The Third Ramsey Numbers for Graphs with at Most Four Edges, *Discrete Mathematics*, **125** (1994) 399-406.
- [-] Yang Yuansheng, see also [SunYJLS, SunYLZ1, SunYLZ2, SunYW, SunYWLX, SunYXL].
- [-] Ye Weiguo, see [SoYL].
- [Yu1]* Yu Song Nian, A Computer Assisted Number Theoretical Construction of $(3, k)$ -Ramsey Graphs, *Annales Universitatis Scientiarum Budapestinensis, Sect. Comput.*, **10** (1989) 35-44.
- [Yu2]* Yu Song Nian, Maximal Triangle-free Circulant Graphs and the Function $K(c)$ (in Chinese), *Journal of Shanghai University, Natural Science*, **2** (1996) 678-682.

Z

- [-] Zang Wenan, see [LiRZ1, LiRZ2, LiTZ, LiZa1, LiZa2].
- [Zeng] Zeng Wei Bin, Ramsey Numbers for Triangles and Graphs of Order Four with No Isolated Vertex, *Journal of Mathematical Research & Exposition*, **6** (1986) 27-32.
- [ZZ1] Zhang Ke Min and Zhang Shu Sheng, Some Tree-Stars Ramsey Numbers, *Proceedings of the Second Asian Mathematical Conference 1995*, 287-291, World Sci. Publishing, River Edge, NJ, 1998.

- [ZZ2] Zhang Ke Min and Zhang Shu Sheng, The Ramsey Numbers for Stars and Stripes, *Acta Mathematica Scientia*, **25A** (2005) 1067-1072.
- [-] Zhang Ke Min, see also [BJYHRZ, ChenZZ1, ChenZZ2, ChenZZ3, ChenZZ4, ChenZZ5, ChenZZ6, HWSYZH, HYZ, HZ1, HZ2, MZ, ShZ1, ShZ2, YHZ1, YHZ2, ZhaCZ1, ZhaCZ2, ZZ3].
- [ZhaCC] Lianmin Zhang, Yaojun Chen and T.C. Edwin Cheng, The Ramsey Numbers for Cycles versus Wheels of Even Order, to appear in the *European Journal of Combinatorics*.
- [-] Zhang Shu Sheng, see [ZZ1, ZZ2].
- [-] Zhang Xiaoxian, see [XieZ].
- [-] Zhang Yuming, see [CLRZ].
- [Zhang1] Zhang Yunqing, On Ramsey Numbers of Short Paths versus Large Wheels, *Ars Combinatoria*, **89** (2008) 11-20.
- [Zhang2] Zhang Yunqing, The Ramsey Numbers for Stars of Odd Small Order versus a Wheel of Order Nine, *Nanjing Daxue Xuebao Shuxue Bannian Kan*, ISSN 0469-5097, **25** (2008) 35-40.
- [ZhaCZ1] Yunqing Zhang, Yaojun Chen and Kemin Zhang, The Ramsey Numbers for Stars of Even Order versus a Wheel of Order Nine, *European Journal of Combinatorics*, **29** (2008) 1744-1754.
- [ZhaCZ2] Yunqing Zhang, Yaojun Chen and Kemin Zhang, The Ramsey Numbers for Trees of High Degree versus a Wheel of Order Nine, *manuscript*, (2009).
- [ZZ3] Yunqing Zhang and Ke Min Zhang, The Ramsey Number $R(C_8, K_8)$, *Discrete Mathematics*, **309** (2009) 1084-1090.
- [-] Zhang Yunqing, see also [ChenCZ1, ChenCZ2, ChenZZ1, ChenZZ2, ChenZZ3, ChenZZ4, ChenZZ5, ChenZZ6, CheCZN].
- [-] Zhang Zhengyou, see [SLZL].
- [ZLLS] Zhang Zhongfu, Liu Linzhong, Li Jinwen and Song En Min, Some Properties of Ramsey Numbers, *Applied Mathematics Letters*, **16** (2003) 1187-1193.
- [-] Zheng Wenping, see [SunYLZ1, SunYLZ2].
- [Zhou1] Zhou Huai Lu, Some Ramsey Numbers for Graphs with Cycles (in Chinese), *Mathematica Applicata*, **6** (1993) 218.
- [Zhou2] Zhou Huai Lu, The Ramsey Number of an Odd Cycle with Respect to a Wheel (in Chinese), *Journal of Mathematics, Shuxue Zazhi* (Wuhan), **15** (1995) 119-120.
- [Zhou3] Zhou Huai Lu, On Book-Wheel Ramsey Number, *Discrete Mathematics*, **224** (2000) 239-249.

Out of 526 references gathered above, 443 appeared in 83 different periodicals, among which most articles were published in: *Discrete Mathematics* 64, *Journal of Combinatorial Theory* (old, Series A and B) 49, *Journal of Graph Theory* 46, *Ars Combinatoria* 27, *Journal of Combinatorial Mathematics and Combinatorial Computing* 20, *Electronic Journal of Combinatorics* 19, and *Utilitas Mathematica* 17. The results of 110 references depend on computer algorithms.